


(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
22 November 2001 (22.11.2001)

PCT

(10) International Publication Number  
**WO 01/87038 A2**

- (51) International Patent Classification: Not classified
- (21) International Application Number: PCT/US01/14357
- (22) International Filing Date: 3 May 2001 (03.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/204,667 17 May 2000 (17.05.2000) US
- (71) Applicant: **ORTHO-MCNEIL PHARMACEUTICAL, INC.** [US/US]; U.S. Route 202, Raritan, NJ 08869 (US).
- (72) Inventors: **SUI, Zhihua**; 11 Running Brook Circle, Flemington, NJ 08892 (US). **MACIELAG, Mark, J.**; 8 Seneca Trail, Banchburg, NJ 08876 (US).
- (74) Agents: **JOHNSON, Philip, S.**; Johnson & Johnson, One Johnson & Johnson Plaza, New Brunswick, NJ 08933 et al. (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title:  $\beta$ -CARBOLINE DERIVATIVES USEFUL AS INHIBITORS OF PHOSPHODIESTERASE

**(57) Abstract:** The present invention relates to novel  $\beta$ -carboline derivatives of the general formula (I), wherein all the variables are as described within the specification, useful as phosphodiesterase inhibitors. The present invention further relates to use of the described derivatives for the treatment of diseases and conditions related to PDE, for example male erectile dysfunction.

WO 01/87038 A2

## $\beta$ -CARBOLINE DERIVATIVES USEFUL AS INHIBITORS OF PHOSPHODIESTERASE

### CROSS REFERENCE TO RELATED APPLICATION

5        This application claims priority from United States provisional application Serial No. 60/204,667 filed May 17, 2000, the contents of which are hereby incorporated by reference.

### FIELD OF THE INVENTION

10        The present invention relates to novel  $\beta$ -carboline derivatives, useful as phosphodiesterase inhibitors. The invention further relates to synthesis of the  $\beta$ -carboline derivatives and intermediates used in their preparation. The present invention additionally relates to use of the described derivatives for the treatment of diseases and  
15        conditions related to PDE, for example male erectile dysfunction.

### BACKGROUND OF THE INVENTION

20        Erectile dysfunction (ED) is defined as the inability to achieve or maintain an erection sufficiently rigid for satisfactory sexual intercourse. Currently it is estimated that approximately 7-8% of the male population suffers from some degree of ED, the equivalent of at least 20 million men in the United States alone. Since the likelihood of ED increases with age, it is projected that the incidence of this condition will rise in the future as the average age of the population increases.

25        Male erectile dysfunction may be the consequence of psychogenic and/or organic factors. Although ED is multi-factorial, certain sub-groups within the male population are more likely to present with the symptoms of the disorder. In particular, patients with diabetes, hypertension, heart disease, and multiple sclerosis have a  
30        particularly high prevalence of ED. In addition, patients who take certain

classes of drugs such as antihypertensives, antidepressants, sedatives, and anxiolytics are more prone to suffer from ED.

Treatments for ED include a variety of pharmacologic agents, vacuum devices, and penile prostheses. Among the pharmacologic agents, papaverine, phentolamine, and alprostadil are currently used in practice. These agents are only effective after direct intracavernosal or intraurethral injection, and are associated with side effects such as priapism, fibrosis, penile pain and hematoma at the injection site. Vacuum devices are a noninvasive alternative treatment for ED. These devices produce an erection by creating a negative pressure around the shaft of the penis resulting in an increased blood flow into the corpus cavernosum via passive arterial dilation. Although this form of therapy is frequently successful in ED of organic origin, complaints include the lack of spontaneity and the time involved in using a mechanical device, and difficulty and discomfort with ejaculation. A variety of semi-rigid or inflatable penile prostheses have been used with some success, particularly in diabetic men. These devices are generally considered when other treatment options have failed, and are associated with an increased risk of infection and ischemia.

Recently, the phosphodiesterase V (PDEV) inhibitor, sildenafil (Viagra®) was approved by the FDA as an orally effective medication for the treatment of ED. Sildenafil, 5-[2-ethoxy-5-(4-methylpiperazin-1-ylsulphonyl)phenyl]-1-methyl-3-n-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one and a number of related analogs and their use as antianginal agents are described in U.S. Patent Nos. 5,250,534 and 5,346,901. The use of sildenafil and related analogs for treating male erectile dysfunction is described in PCT International Application Publication No. WO 94/28902, published December 22, 1994. In clinical studies, the drug improved sexual function in about 70% of the men who suffer from ED of psychogenic or organic etiology. However, the drug showed less dramatic efficacy in patients who had undergone a radical prostatectomy, with improved erections in 43% of patients who took sildenafil versus 15% on placebo. In addition, the use of sildenafil is associated with several undesirable side effects including headache, flushing and

disrupted color vision which result from non-selective effects on a variety of tissues. In spite of these shortcomings, the drug is viewed by patients as preferable to other treatments which involve the introduction of medication directly into the penis via injection, the use of an external device or a surgical procedure.

5

Daugan et.al, in US Patent No. 5,859,009 (EP 0740668 B1 and WO9519978) describe the synthesis of tetracyclic derivatives as inhibitors of cyclic guanosine 3',5'-monophosphate specifically phosphodiesterase, and their use in treating cardiovascular disorders. Daugan et.al., in WO97/03675 teach the use of tetracyclic derivatives for the treatment of impotence.

10

Bombrun et al., in WO 97/43287 describe a series of carboline derivatives, more specifically 2-(substituted alkyl carbonyl) substituted carboline derivatives and their use in treating cardiovascular disorders as inhibitors of cyclic guanosine 3,5-monophosphate, specifically phosphodiesterase.

15

Ellis et. al., in WO 94/28902 and EP0702555 B1 describes a series of pyrazolpyrimidinone derivatives and their use in treating erectile dysfunction. Campbell, S. F. in WO96/16657 teaches the use of bicyclic heterocyclic compounds for the treatment of impotence (pyrazolopyrimidones); while Campbell et al, in WO,96/16644 teach the use of selective cGMP PDE inhibitors for the treatment of erectile dysfunction.

20

Ohashi et al., in WO9745427 disclose tetracyclic pyridocarbazole derivatives having cGMP PDE inhibitory effects.

25

Fourtillan et. al., in WO 96/08490 A1 describe a series of carboline derivatives and their use in the treatment of diseases associated with melatonin activity disorders. Ueki et. al., in US Patent No 5,126,448 describe pyridine and 1,2,3,4-tetrahydropyridine derivatives useful as psychotropic drugs having antianxiety effects. Atkinson et. al., in US Patent No. 3,328,412 describe 1-aryl- and 1-

30

heteroaryl-2-acyl-1,2,3,4-tetrahydro- $\beta$ -carboline derivatives having long lasting analgesic properties.

Sexually stimulated penile erection results from a complex interplay of physiological processes involving the central nervous system, the peripheral nervous system, and the smooth muscle. Specifically, release of nitric oxide from the non-adrenergic, non-cholinergic nerves and endothelium activates guanylyl cyclase and increases intracellular cGMP levels within the corpus cavernosum. The increase in intracellular cGMP reduces intracellular calcium levels, resulting in trabecular smooth muscle relaxation, which, in turn, results in corporal volume expansion and compression of the sub-tunical venules leading to penile erection.

PDEV has been found in human platelets and vascular smooth muscle, suggesting a role for this enzyme in the regulation of intracellular concentrations of cGMP in cardiovascular tissue. In fact, inhibitors of PDEV have been shown to produce endothelial-dependent vasorelaxation by potentiating the increases in intracellular cGMP induced by nitric oxide. Moreover, PDEV inhibitors selectively lower the pulmonary arterial pressure in animal models of congestive heart failure and pulmonary hypertension. Hence in addition to their utility in ED, PDEV inhibitors would likely be of therapeutic benefit in conditions like heart failure, pulmonary hypertension, and angina.

Agents that increase the concentration of cGMP in penile tissue, either through enhanced release or reduced breakdown of cGMP, are expected to be effective treatments for ED. The intracellular levels of cGMP are regulated by the enzymes involved in its formation and degradation, namely the guanylate cyclases and the cyclic nucleotide phosphodiesterases (PDEs). To date, at least nine families of mammalian PDEs have been described, five of which are capable of hydrolyzing the active, cGMP, to the inactive, GMP, under physiological conditions (PDEs I, II, V, VI, and IX). PDE V is the predominant isoform in human corpus cavernosum. Inhibitors of PDEV, therefore, would be expected to increase the concentration of

cGMP in the corpus cavernosum and enhance the duration and frequency of penile erection.

Additionally, selective PDE inhibitors are known to be useful in the treatment  
5 of various disorders and conditions including male erectile dysfunction (ED), female sexual arousal dysfunction, female sexual dysfunction related to blood flow and nitric oxide production in the tissues of the vagina and clitoris, premature labor, dysmenorrhea, cardiovascular disorders, atherosclerosis, arterial occlusive disorders, thrombosis, coronary rest stenosis, angina pectoris, myocardial  
10 infarction, heart failure, ischemic heart disorders, hypertension, pulmonary hypertension, asthma, intermittent claudication and diabetic complications.


Accordingly, it is an object of the invention to identify compounds which increase the concentration of cGMP in penile tissue through the inhibition of  
15 phosphodiesterases, specifically PDEV. It is another object of the invention to identify compounds which are useful for the treatment of sexual dysfunction, particularly erectile dysfunction and/or impotence in male animals and sexual dysfunction in female animals. Still another object of the invention is to identify methods for treating sexual dysfunction, especially erectile dysfunction, using the compounds of the present  
20 invention.

It is another object of the invention to identify compounds which are useful for the treatment of conditions of disorders mediated by PDEV, such as male erectile dysfunction, female sexual dysfunction, cardiovascular disorders,  
25 atherosclerosis, arterial occlusive disorders, thrombosis, coronary rest stenosis, angina pectoris, myocardial infarction, heart failure, ischemic heart disorders, hypertension, pulmonary hypertension, asthma, intermittent claudication or diabetic complications.

30 We now describe a series of  $\beta$ -carboline derivatives with the ability to inhibit phosphodiesterase type V in enzyme assays and increase the concentration of cGMP in cavernosal tissue in vitro.

### SUMMARY OF THE INVENTION

The present invention provides novel  $\beta$ -carboline derivative compounds  
 5 useful as phosphodiesterase inhibitors. More particularly, the present invention is directed to compounds of the general formula (I):



wherein

$R^1$  is independently selected from the group consisting of halogen, nitro,  
 10 hydroxy,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy,  $-NH_2$ ,  $-NHR^A$ ,  $-N(R^A)_2$ ,  $-OR^A$ ,  $-C(O)NH_2$ ,  
 $-C(O)NHR^A$ ,  $-C(O)N(R^A)_2$ ,  $-NC(O)-R^A$ ,  $-SO_2NHR^A$ ,  $-SO_2N(R^A)_2$ , phenyl (optionally  
 substituted with 1 to 3  $R^B$ ) and heteroaryl (optionally substituted with 1 to 3  $R^B$ );

where each  $R^A$  is independently is independently selected from the group  
 consisting of  $C_1$ - $C_8$ alkyl, aryl (optionally substituted with 1 to 3  $R^B$ ),  $C_1$ - $C_8$ aralkyl  
 15 (optionally substituted with 1 to 3  $R^B$ ) and heteroaryl (optionally substituted with 1 to  
 3  $R^B$ );

where each  $R^B$  is independently selected from the group consisting of  
 halogen, hydroxy, nitro, cyano,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy,  $C_1$ - $C_8$ alkoxycarbonyl,  
 carboxy $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkylsulfonyl, trifluoromethyl, trifluoromethoxy, amino,  
 20 acetylamino, di( $C_1$ - $C_8$ alkyl)amino, di( $C_1$ - $C_8$ alkyl)amino $C_1$ - $C_8$ alkoxy, di( $C_1$ -  
 $C_8$ alkyl)aminoacetyl $C_1$ - $C_8$ alkyl, di( $C_1$ - $C_8$ alkyl)aminoacetylamino, carboxy $C_1$ -  
 $C_8$ alkylcarbonylamino, hydroxy $C_1$ - $C_8$ alkylamino,  $NHR^A$ ,  $N(R^A)_2$  and  
 heterocycloalkyl $C_1$ - $C_8$ alkoxy;

$n$  is an integer from 0 to 4;

25  $X$  is selected from the group consisting of O, S and  $NR^D$ ;

where  $R^D$  is selected from the group consisting of hydrogen, hydroxy,  $-OR^A$ ,  $C_1$ - $C_8$ alkyl (wherein the alkyl is optionally substituted with one to three substituent independently selected from halogen, carboxy, amino,  $C_1$ - $C_8$ alkylamino,  $di(C_1$ - $C_8$ alkyl)amino,  $C_1$ - $C_8$ alkoxycarbonyl, heteroaryl or heterocycloalkyl), heteroaryl and  
 5 heteroarylcarbonyl (wherein the heteroaryl may be optionally substituted with phenyl or substituted phenyl, where the phenyl substituents are one to three  $R^B$ );

$R^2$  is selected from the group consisting of  $C_5$ - $C_{10}$ alkyl (optionally substituted with 1 to 3  $R^C$ ), aryl (optionally substituted with 1 to 3  $R^B$ ), heteroaryl (optionally substituted with 1 to 3  $R^B$ ) and heterocycloalkyl (optionally substituted with 1 to 3  
 10  $R^B$ );

where each  $R^C$  is independently selected from the group consisting of halogen, hydroxy, nitro,  $NH_2$ ,  $NHR^A$  and  $N(R^A)_2$ ;


Z is selected from the group consisting of  $CH_2$ ,  $CHOH$  and  $C(O)$ ; provided that when Z is  $CHOH$  or  $C(O)$ , then X is  $NH$ ;

15  $R^4$  is selected from the group consisting of hydrogen, hydroxy, carboxy,  $C_1$ - $C_6$ alkylcarbonyl,  $C_1$ - $C_6$ alkoxycarbonyl,  $di(C_1$ - $C_8$ alkyl)aminoalkoxycarbonyl,  $di(C_1$ - $C_8$ alkyl)amino $C_1$ - $C_8$ alkylaminocarbonyl, and  $-COR^F$ ;

where  $R^F$  is selected from the group consisting of  $C_1$ - $C_8$ alkyl,  $NH_2$ ,  $NHR^A$ ,  $NR^A_2$ ,  $-C_1$ - $C_8$ alkyl- $NH_2$ ,  $-C_1$ - $C_8$ alkyl- $NHR^A$ ,  $-C_1$ - $C_8$ alkyl- $NR^A_2$  and  $-NH$ - $C_1$ - $C_8$ alkyl-  
 20  $NR^A_2$ ;

a is an integer from 0 to 1;

Y is selected from the group consisting of  $CH_2$ ,  $C(O)$ ,  $C(O)O$ ,  $C(O)-NH$  and  $SO_2$ ;





is selected from the group consisting of naphthyl, heteroaryl and  
 25 heterocycloalkyl;

m is an integer from 0 to 2;

$R^3$  is independently selected from the group consisting of halogen, nitro,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy, trifluoromethyl, trifluoromethoxy, phenyl (optionally substituted with 1 to 3  $R^B$ ), phenylsulfonyl, naphthyl,  $C_1$ - $C_8$ alkyl, heteroaryl (optionally substituted with 1 to 3  $R^B$ ),  $NH_2$ ,  $NHR^A$  and  $N(R^A)_2$ ;  
 30





provided that when  is 2-furyl or 2-thienyl, then m is an integer from 1 to 2;

and pharmaceutically acceptable salts thereof.

5 Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. An illustration of the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier.

10 Illustrating the invention is a process for making a pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

Exemplifying the invention are methods of treating sexual dysfunction, for example, male erectile dysfunction, impotence, female sexual dysfunction, for  
15 example female sexual arousal dysfunction, female sexual dysfunction related to blood flow and nitric oxide production in the tissues of the vagina and clitoris, premature labor and / or dysmenorrhea in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

20

An example of the invention is a method for increasing the concentration of cGMP in penile tissue through the inhibition of phosphodiesterases, specifically PDEV, in a male subject in need thereof comprising administering to the subject an  
25 effective amount of any of the compounds or pharmaceutical compositions described above.

Further exemplifying the invention is a method of producing endothelial-dependent vasorelaxation by potentiating the increases in intracellular cGMP induced by nitric oxide in a subject in need thereof comprising administering to the

subject an effective amount of any of the compounds or pharmaceutical compositions described above.

Further illustrating the invention is a method of treating a condition selected from the group consisting of male erectile dysfunction (ED), impotence, female sexual dysfunction, female sexual arousal dysfunction, female sexual dysfunction related to blood flow and nitric oxide production in the tissues of the vagina and clitoris, premature labor, dysmenorrhea, cardiovascular disorders, atherosclerosis, arterial occlusive disorders, thrombosis, coronary rest stenosis, angina pectoris, myocardial infarction, heart failure, ischemic heart disorders, hypertension, pulmonary hypertension, asthma, intermittent claudication and diabetic complications in a subject in need thereof comprising administering to the subject a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

15

An example of the invention is the use of any of the compounds described above in the preparation of a medicament for: (a) treating sexual dysfunction, especially male erectile dysfunction, (b) treating impotence, (c) increasing the concentration of cGMP in penile tissue through inhibition of phosphodiesterase, especially PDEV and/or (d) treating a condition selected from the group consisting of premature labor, dysmenorrhea, cardiovascular disorders, atherosclerosis, arterial occlusive disorders, thrombosis, coronary rest stenosis, angina pectoris, myocardial infarction, heart failure, ischemic heart disorders, hypertension, pulmonary hypertension, asthma, intermittent claudication and diabetic complications in a subject in need thereof.

25

#### DETAILED DESCRIPTION OF THE INVENTION


The present invention provides novel  $\beta$ -carboline derivatives useful for the treatment of sexual dysfunction, particularly male erectile dysfunction (ED). Although the compounds of the present invention are useful primarily for the treatment of male sexual dysfunction or erectile dysfunction, they may also be useful

30

for the treatment of female sexual dysfunction, for example female sexual arousal dysfunction, female sexual dysfunction related to blood flow and nitric oxide production in the tissue of the vagina and clitoris, and of premature labor and dysmenorrhea.

5

More particularly, the compounds of the present invention are of the formula (I):



wherein all variables are as defined above.

10

Preferably,  $n$  is 0. Preferably,  $m$  is an integer from 0 to 1.

In an embodiment of the present invention  $X$  is selected from  $S$  or  $NR^D$ , wherein  $R^D$  is selected from the group consisting of hydrogen, halo $C_1$ - $C_6$ alkyl, di( $C_1$ - $C_4$ alkyl)amino $C_1$ - $C_6$ alkyl, heteroaryl, heteroaryl $C_1$ - $C_4$ alkyl, heterocycloalkyl $C_1$ - $C_4$ alkyl, carboxy $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxycarbonyl $C_1$ - $C_4$ alkyl and heteroarylcarbonyl; wherein the heteroaryl is further optionally substituted with phenyl or substituted phenyl, wherein the substituents on the phenyl are one to two independently selected from  $R^B$ ; and wherein each  $R^B$  is independently selected from the group consisting of halogen, nitro,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, trifluoromethyl, trifluoromethoxy, amino and di( $C_1$ - $C_4$ alkyl)amino. Preferably,  $X$  is selected from  $S$  or  $NR^D$ , where  $R^D$  is selected from the group consisting of hydrogen, di(methyl)aminoethyl, di(methyl)amino- $n$ -propyl, di(ethyl)aminoethyl, di(ethyl)amino- $n$ -butyl,  $N$ -pyrrolidinyethyl,  $N$ -morpholinylethyl, 2-pyridylmethyl, 4-pyridylmethyl, 5-(4-methylphenyl)-2-pyrimidinyl, carboxymethyl, carboxyethyl, 4-chloro- $n$ -butyl, 2-(5-(3-trifluoromethylphenyl)furyl)carbonyl, 2-(5-(3-nitrophenyl)furyl)carbonyl,




25


methoxycarbonylmethyl, methoxycarbonylethyl and 2-benzoxazolyl. More preferably, X is  $\text{NR}^{\text{D}}$ , where  $\text{R}^{\text{D}}$  is selected from the group consisting of hydrogen, di(methyl)aminoethyl, 4-pyridylmethyl, 2-pyridylmethyl, N-morpholinylethyl, carboxyethyl, carboxymethyl, di(ethyl)aminoethyl, N-pyrrolidinylethyl and 5-(4-methylphenyl)-2-pyrimidinyl. Most preferably, X is  $\text{NR}^{\text{D}}$ , where  $\text{R}^{\text{D}}$  is selected from the group consisting of hydrogen, di(methyl)aminoethyl, N-morpholinylethyl, carboxymethyl and N-pyrrolidinylethyl;


Preferably, Z is selected from the group consisting of  $\text{CH}_2$  and  $\text{C(O)}$ ; provided that when Z is  $\text{C(O)}$ , then X is NH.

Preferably, Y is selected from the group consisting of  $\text{C(O)}$ ,  $\text{SO}_2$  and  $\text{CH}_2$ . More preferably, Y is selected from the group consisting of  $\text{C(O)}$  and  $\text{CH}_2$ . Most preferably, Y is  $\text{C(O)}$ .

15

In an embodiment of the present invention  is selected from the group consisting of naphthyl and heteroaryl. Preferably,  is selected from the group consisting of naphthyl, 2-pyrimidinyl, 2-furyl, 3-furyl, 2-benzofuryl, 2-thienyl, 2-benzothienyl, 2-benzothiazolyl, 2-benzoxazolyl, 2-benzimidazolyl, 4-thiazolyl, 2-thiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 3-(1,2,5-triazolyl), 4-isoxazolyl, 2-pyridyl and 3-pyridyl. More preferably,  is selected from the group consisting of naphthyl, 2-pyrimidinyl, 2-furyl, 2-benzofuryl, 2-thienyl, 2-benzothienyl, 2-benzothiazolyl, 2-benzoxazolyl, 2-thiazolyl, 4-thiazolyl and 2-pyridyl.



Most preferably,  is selected from the group consisting of 2-pyrimidinyl, 2-furyl, 2-benzofuryl, 2-benzoxazolyl, 2-thiazolyl and 2-pyridyl.

In an embodiment of the present invention,  $R^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl, 3,4-(difluoro)methylenedioxyphenyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzo-[1,4]-dioxin-6-yl, pyridyl, phenyl and substituted phenyl; wherein the phenyl substituents are one to two substituents independently selected from halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, trifluoromethyl, cyano, nitro,  $C_1$ - $C_4$ alkoxycarbonyl,  $di(C_1-C_4alkyl)amino$  or  $di(C_1-C_4alkyl)aminoC_1-C_4alkoxy$ .

10 Preferably,  $R^2$  is selected from the group consisting of phenyl, 3,4-methylenedioxyphenyl, 3,4-(difluoro)methylenedioxyphenyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzo-[1,4]-dioxin-6-yl, 4-pyridyl, 3-pyridyl, 4-cyanophenyl, 3-nitrophenyl, 4-nitrophenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethyl-4-chlorophenyl, 3,4-

15 dichlorophenyl, 4-chlorophenyl, 4-methoxycarbonylphenyl, 3,4-dimethoxyphenyl, 4-(dimethylamino)phenyl and 4-(N-(3-dimethylamino)-n-propoxy)phenyl. More preferably,  $R^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl, 2,3-dihydrobenzofuryl and 2,3-dihydrobenzo-[1,4]-dioxin-6-yl. Most preferably,  $R^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl and 2,3-

20 dihydrobenzofuryl.

Preferably,  $R^4$  is selected from the group consisting of hydrogen, carboxy,  $C_1$ - $C_4$ alkoxycarbonyl,  $di(C_1-C_4alkyl)aminoC_1-C_4alkoxycarbonyl$  and  $di(C_1-C_4alkyl)aminoC_1-C_4alkylaminocarbonyl$ . More preferably,  $R^4$  is selected from the

25 group consisting of hydrogen, carboxy, dimethylaminoethoxycarbonyl, dimethylaminoethylaminocarbonyl and methoxycarbonyl. Most preferably,  $R^4$  is hydrogen.

In an embodiment of the present invention,  $R^3$  is independently selected from

30 the group consisting of halogen, nitro,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, trifluoromethyl,  $C_1$ -

C<sub>4</sub>alkyl, pyrazinyl, pyridyl, halogen substituted pyridyl, dimethyl substituted imidazolyl, phenyl, phenylsulfonyl and substituted phenyl; wherein the substituents on the phenyl are one or more substituents independently selected from halogen, hydroxy, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, trifluoromethyl, trifluoromethoxy, nitro, amino, acetylamino, C<sub>1</sub>-C<sub>4</sub>alkylsulfonyl, carboxyC<sub>1</sub>-C<sub>4</sub>alkylcarbonylamino, hydroxyC<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)aminoC<sub>1</sub>-C<sub>4</sub>alkoxy, di(C<sub>1</sub>-C<sub>4</sub>alkyl)aminoacetylamino or heterocycloalkylC<sub>1</sub>-C<sub>4</sub>alkoxy. Preferably, R<sup>3</sup> is independently selected from the group consisting of chloro, bromo, methyl, n-propyl, t-butyl, methoxy, trifluoromethyl, nitro, phenyl, benzyl, phenylsulfonyl, 4-hydroxyphenyl, 4-chlorophenyl, 4-methylphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 5-trifluoromethylphenyl, 4-methoxyphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 4-aminophenyl, 2-nitro-4-chlorophenyl, 2-nitro-4-methylphenyl, 2-nitro-4-methylsulfonylphenyl, 3-acetylaminophenyl, 4-acetylaminophenyl, 4-(3-carboxy-n-propyl)carbonylaminophenyl, 2-chloro-5-trifluoromethylphenyl, 4-(4-hydroxy-n-butyl)aminophenyl, 2-(dimethylamino)acetylaminophenyl, 4-[2-(N-pyrrolidinyl)ethoxy]phenyl, 4-[2-(4-morpholinyl)ethoxy]phenyl, 4-(2-(dimethylamino)ethoxy)phenyl, 4-pyrazinyl, 2,3-dimethyl-3H-imidazolyl, 2-pyridyl and 3-pyridyl. More preferably, R<sup>3</sup> is selected from the group consisting of bromo, t-butyl, methoxy, trifluoromethyl, nitro, phenyl, 4-chlorophenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 4-methylphenyl, 4-methoxyphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 2-nitro-4-chlorophenyl, 2-nitro-4-methylphenyl, 2-nitro-4-methylsulfonylphenyl, 4-(3-carboxy-n-propyl)carbonylaminophenyl, 2-chloro-5-trifluoromethylphenyl, 4-(4-hydroxy-n-butyl)aminophenyl, 2-(dimethylamino)acetylaminophenyl, 4-pyrazinyl, 2-pyridyl and 2,3-dimethyl-3H-imidazol-4-yl. Most preferably, R<sup>3</sup> is selected from the group consisting of t-butyl, methoxy, nitro, phenyl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 2-nitro-4-methylsulfonylphenyl, 2-(dimethylamino)acetylaminophenyl, 2-pyridyl and 2,3-dimethyl-3H-imidazol-4-yl.

For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the

preparation of compounds according to this invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as

5 hydrochloric acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium

10 salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include the following:

acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate,

15 gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, pamoate (embonate), palmitate, pantothenate,

20 phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate.

The present invention includes within its scope prodrugs of the compounds of this invention. In general, such prodrugs will be functional derivatives of the

25 compounds which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various disorders described with the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the

30 patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

As used herein, unless otherwise noted, "halogen" shall mean chlorine, bromine, fluorine and iodine.

The term "alkyl", whether used alone or as part of a substituent group, shall mean straight or branched chain alkanes of one to ten carbon atoms, or any number within this range. For example, alkyl radicals include, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, n-hexyl and 2-methylpentyl.

The term "alkoxy" shall denote an oxygen ether radical of the above described straight or branched chain alkyl group. For example, alkoxy radicals include methoxy, ethoxy, n-propoxy, n-butoxy, sec-butoxy, tert-butoxy, and the like.

The term "aryl" indicates an aromatic groups such as phenyl, naphthyl, and the like.

The term "aralkyl" denotes an alkyl group substituted with an aryl group. For example, benzyl, phenylethyl, and the like.



The term "heteroaryl" as used herein represents a stable five or six membered monocyclic aromatic ring system containing one to three heteroatoms independently selected from N, O or S; and any nine or ten membered bicyclic aromatic ring system containing carbon atoms and one to four heteroatoms independently selected from N, O or S. The heteroaryl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of heteroaryl groups include, but are not limited to pyridyl, pyrimidinyl, thienyl, furyl, imidazolyl, isoxazolyl, oxazolyl, pyrazolyl, pyrazinyl, pyrrolyl, thiazolyl, thiadiazolyl, triazolyl, benzimidazolyl, benzofuranyl, benzothienyl, benzisoxazolyl, benzoxazolyl, indazolyl, indolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl, quinolinyl or isoquinolinyl. Particularly preferred heteroaryl groups include pyridyl, pyrazolyl, furyl, thiazolyl, thienyl, imidazolyl, isoxazolyl, pyrazinyl, pyrimidinyl, triazolyl, benzofuryl, benzothienyl, benzimidazolyl, benzothiazolyl and benzoxazolyl.


The term "cycloalkyl" as used herein represents a stable three to eight membered monocyclic ring structure consisting of saturated carbon atoms. Suitable examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

The term "heterocycloalkyl" represents a stable saturated or partially unsaturated, three to eight membered monocyclic ring structure containing carbon atoms and one to four, preferably one to two, heteroatoms independently selected from N, O or S; and any stable saturated, partially unsaturated or partially aromatic, nine to ten membered bicyclic ring system containing carbon atoms and one to four heteroatoms independently selected from N, O or S. The heterocycloalkyl may be attached at any carbon atom or heteroatom which results in the creation of a stable structure. Suitable examples of heterocycloalkyl groups include pyrrolidinyl, pyrazolidinyl, piperidinyl, piperazinyl, morpholinyl, dithianyl, trithianyl, dioxolanyl, dioxanyl, thiomorpholinyl, 3,4-methylenedioxypyphenyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzo-[1,4]-dioxin-6-yl, 2,3-dihydro-furo[2,3-b]pyridyl, 1,2-(methylenedioxy)cyclohexane, and the like. Particularly preferred heterocycloalkyl

groups include pyrrolidinyl, morpholinyl, 3,4-methylenedioxyphenyl, 2,3-dihydrobenzofuryl and 2,3-dihydrobenzo-[1,4]-dioxin-6-yl.

As used herein, the notation "\*" shall denote the presence of a stereogenic  
5 center.

Under standard nomenclature used throughout this disclosure, the terminal  
portion of the designated side chain is described first, followed by the adjacent  
functionality toward the point of attachment. Thus, for example, a "phenylC<sub>1</sub>-C<sub>6</sub>  
10 alkylaminocarbonylC<sub>1</sub>-C<sub>6</sub>alkyl" substituent refers to a group of the formula



It is intended that the definition of any substituent or variable at a particular  
location in a molecule be independent of its definitions elsewhere in that molecule.  
15 It is understood that substituents and substitution patterns on the compounds of this  
invention can be selected by one of ordinary skill in the art to provide compounds  
that are chemically stable and that can be readily synthesized by techniques known  
in the art as well as those methods set forth herein. It is further intended that when  
n or m is >1, the corresponding R<sup>1</sup> or R<sup>3</sup> substituents may be the same or different.

20

The term "sexual dysfunction" as used herein, includes male sexual  
dysfunction, male erectile dysfunction, impotence, female sexual dysfunction,  
female sexual arousal dysfunction and female sexual dysfunction related to blood  
flow and nitric oxide production in the tissues of the vagina and clitoris.

25

The term "subject" as used herein, refers to an animal, preferably a mammal,  
most preferably a human, who has been the object of treatment, observation or  
experiment.

The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

Abbreviations used in the specification, particularly the Reactions and Examples, are as follows:


Cmpd #	=	Compound ID Number
DCC	=	1,3-Dicyclohexylcarbodiimide
DCM	=	Dichloromethane
DDQ	=	Dichlorodicyanoquinone
DIC	=	Diisopropylcarbodiimide
DIPEA	=	Diisopropylethylamine
DMF	=	N,N-Dimethylformamide
DMSO	=	Dimethylsulfoxide
dppp	=	1,3-Bis(diphenylphosphino) propane
EDTA	=	Ethylenedinitrilotetracetic acid
Fmoc	=	9-Fluorenylmethoxycarbonyl
Fmoc-NCS	=	9-fluorenylmethoxycarbonyl isothiocyanate
HEPES	=	2-[4-(2-Hydroxyethyl)-piperazinyl]- ethanesulfonic acid
LAH	=	Lithium Aluminum Hydride
PDE	=	Phosphodiesterase


$\text{Pd}_2\text{dba}_3$	=	Tris(dibenzylidene acetone) dipalladium (0)
$\text{Pd}(\text{OAc})_2$	=	Palladium (II) Acetate
$\text{Pd}(\text{PPh}_3)_4$	=	Palladium tetrakis(triphenyl phosphine)
Ph	=	Phenyl
PMSF	=	Phenylmethane sulfonyl fluoride
$\text{PPh}_3$	=	Triphenyl phosphine
PyBop	=	(1-Hydroxy-1H-benzotriazolato-O)tri- 1-pyrrolidinyl phosphorous
PyBrop	=	Bromo-tri-1-pyridinyl phosphorous
SNP	=	Sodium nitroprusside
TEA	=	Triethylamine
TFA	=	Trifluoroacetic acid
THF	=	Tetrahydrofuran
TsOH	=	Tosic Acid

Compounds of formula (I) may be prepared according to the processes outlined in more detail below.

5

Compounds of formula (I) wherein  $(Y)_a$  is  $\text{C}(\text{O})$  may be prepared according to a process as outlined in Scheme 1.







SCHEME 1

More particularly, a compound of formula (II), wherein X is O, S or NH, a  
 5 known compound or compound produced by known methods, is reacted with a  
 suitably substituted aldehyde of formula (III), in an organic solvent such as DCM,  
 THF, toluene, and the like, in the presence of an acid catalyst such as TFA, tosic  
 acid, and the like, to produce the corresponding tricyclic compound of formula (IV).

The compound of formula (IV) is reacted with a suitably substituted  
 10 compound of formula (V), wherein A is halogen, in the presence of a base such as  
 triethylamine (TEA), diisopropylethylamine (DIPEA), sodium carbonate and the like,  
 in an organic solvent such as dichloromethane (DCM), N,N'-dimethylformamide  
 (DMF), tetrahydrofuran (THF), and the like; or with a suitably substituted compound  
 of formula (V), wherein A is hydroxy, in the presence of a coupling agent such as  
 15 DCC, DIC, PyBop, PyBrop, and the like, in an organic solvent such as  
 dichloromethane (DCM), N,N'-dimethylformamide (DMF), tetrahydrofuran (THF), and  
 the like; to produce the corresponding compound of formula (Ia).

Alternatively, compounds of formula (I), wherein X is O, S, or NH and (Y)<sub>a</sub> is  
 20 C(O) may be prepared according to a process as outlined in Scheme 2.






SCHEME 2

5 More particularly, a compound of formula (II), wherein  $X$  is O, S or NH, is reacted with a suitably substituted compound of formula (VI), wherein  $A$  is halogen or hydroxy, in an organic solvent such as DCM, THF, DMF, and the like, to produce the corresponding compound of formula (VII).

10 The compound of formula (VII) is cyclized by treatment with  $POCl_3$ , in an organic solvent such as toluene, benzene, and the like, followed by reduction with  $NaBH_4$ , in an organic solvent such as ethanol, isopropanol, and the like, to produce the corresponding compound of formula (IV).

15 The compound of formula (IV) is then reacted with a suitably substituted compound of formula (V) to produce the compound of formula (Ia) as outlined in Scheme 1.


Compounds of formula (I), wherein  $X$  is O, S or NH and  $(Y)_a$  is  $SO_2$  may be prepared according to a process as outlined in Scheme 3.



5        Accordingly, a suitably substituted compound of formula (IV) is reacted with a  
 suitably substituted compound of formula (VIII), wherein A is halogen or hydroxy, a  
 known compound or compound prepared by known methods, in an organic solvent  
 such as DCM, chloroform, DMF, THF, and the like, to produce the corresponding  
 compound of formula (Ib).

10


Compounds of formula (I) wherein X is O, S or NH and  $(Y)_a$  is  $CH_2$  may be  
 prepared according to a process as outlined in Scheme 4.



15

Accordingly, a suitably substituted compound of formula (Ia) is treated with a reducing agent such as LAH, diboron, and the like, preferably LAH, in an organic solvent such as methanol, THF, diethyl ether, and the like, preferably at a temperature in the range of about  $-20$  to  $40^{\circ}\text{C}$ , to produce the corresponding compound of formula (Ic).


Compounds of formula (I) wherein  $(Y)_a$  is  $\text{CH}_2$  and X is NH, may alternatively be prepared according to a process as outlined in Scheme 5.




SCHEME 5

Accordingly, a compound of formula (IVa) is reacted with a suitably substituted compound of formula (IX), wherein Q is halogen, O-tosylate or O-mesolate, in an organic solvent such as DCM, THF, and the like, to produce the corresponding compound of formula (Id).

Compounds of formula (I), wherein X is O, S or NH and  $(Y)_a$  is  $(Y)_0$  (i.e. wherein a is 0, such that Y is absent), may be prepared according to a process as outlined in Scheme 6.








SCHEME 6

More specifically, a compound of formula (IV), a known compound or  
 5 compound produced by known methods, is reacted with a suitably substituted halide  
 of formula (X), a known compound or compound prepared by known methods, in an  
 organic solvent such as toluene, DMF, 1-methyl-2-pyrrolidinone, and the like,  
 preferably at a temperature in the range of about 80 to 250°C, to produce the  
 corresponding compound of formula (Ie).

10

Compounds of formula (I) wherein X is  $\text{NR}^{\text{D}}$  may be prepared by according to  
 a process as outlined in Scheme 7.




SCHEME 7

15


Accordingly, a compound of formula (If) is reacted with a compound of  
 formula (XI), wherein Z is halogen, hydroxy, O-toylate or O-mesolate and a base  
 such as sodium hydride, potassium t-butoxide, and the like, in a solvent such DMF,  
 20 1-methyl-2-pyrrolidinone, and the like, to produce the corresponding compound of  
 formula (Ig).


Compounds of formula (I) wherein Z is CH-OH or C(O) may be prepared according to a process as outlined in Scheme 8.



SCHEME 8

More particularly, a compound of formula (If) is treated with an oxidizing agent such as DDQ, chloranil, and the like, in a solvent such as THF, methanol, water, and the like, preferably at a temperature in the range of about  $-78$  to about  $30^\circ\text{C}$ , to produce a mixture of the corresponding compounds of formula (Ih) and (Ii). Preferably, the compounds of formula (Ih) and (Ii) are separated by known methods, such as recrystallization, column chromatography, and the like.

Compounds of formula (I) wherein  is 2-thiazolyl, may be prepared according to a process as outlined in Scheme 9.



5

SCHEME 9


Accordingly, a suitably substituted compound of formula (IVa) is reacted with Fmoc-NCS, in an organic solvent such as DCM, DMF, THF, and the like, preferably at room temperature, to produce the corresponding compound of formula (XII).

10 The compound of (XII) is reacted with 20% piperidine, in an alcohol such as methanol, ethanol, and the like, to produce the corresponding amine of formula (XIII).

The amine of formula (XIII) is reacted with a suitably substituted  $\alpha$ -halo methyl ketone of formula (XIV), in the presence of an organic solvent or mixture  
 15 such as DMF, ethanol:dioxane, and the like, in the presence of a base such as TEA,

DIPEA, and the like, preferably at a temperature of about 70°C, to produce the corresponding compound of formula (lj).


Compounds of formula (I) wherein (Y)<sub>a</sub> is C(O)O, may be prepared according  
5 to the process outlined in Scheme 10.



SCHEME 10

10 More particularly, a compound of formula (IV) is reacted with a suitably substituted chloroformate of formula (XV) or an anhydride of formula (XVI) in an organic solvent such as DCM, DMF, THF, and the like, to produce the corresponding compound of formula (Ik).

15 Compounds of formula (I) wherein (Y)<sub>a</sub> is C(O)-NH may be prepared according to a process as outlined in Scheme 11.



SCHEME 11

5           Accordingly, a compound of formula (IV) is reacted with a suitably substituted compound of formula (XVII), in an organic solvent such as DCM, DMF, THF, and the like, to produce the corresponding compound of formula (Im).

10           Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the

15           formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be

20           resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

10       The utility of the compounds to treat sexual dysfunction can be determined according to the procedures described in Example 10, 11 and 12 herein.

15       The present invention therefore provides a method of treating sexual dysfunction in a subject in need thereof, which comprises administering any of the compounds as defined herein in a quantity effective to treat sexual dysfunction. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral. The quantity of the compound which is effective for treating sexual dysfunction is between 0.1 mg per kg and 20 mg per kg of subject body weight.

25       The present invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a

pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present  
5 invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type  
10 described above containing from 1 to about 1000 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The  
15 two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

20

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as  
25 elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

30

The method of treating sexual dysfunction, particularly male erectile dysfunction (ED) described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a

pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 1 mg and 1000 mg, preferably about 10 to 500 mg, of the compound, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not  
5 limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixers, emulsions, and suspensions.  
10 Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of  
15 two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.  
20

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can  
25 also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar,  
30 bentonite, xanthan gum and the like.



The liquid forms may include suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are  
5 employed when intravenous administration is desired.

The compound of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of  
10 phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble  
15 polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxyethylaspartamidephenol, or polyethyl-eneoxidepolylysine substituted with palmitoyl residue. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of  
20 a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Compounds of this invention may be administered in any of the foregoing  
25 compositions and according to dosage regimens established in the art whenever treatment of sexual dysfunction, particularly male erectile dysfunction (ED) is required.

The daily dosage of the products may be varied over a wide range from 1 to 1,000 mg per adult human per day. For oral administration, the compositions are  
30 preferably provided in the form of tablets containing, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily

supplied at a dosage level of from about 0.1 mg/kg to about 20 mg/kg of body weight per day. Preferably, the range is from about 0.2 mg/kg to about 10 mg/kg of body weight per day, and especially from about 0.5 mg/kg to about 10 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 4 times per day.

5

Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of administration, will result in the need to adjust dosages.

The following Examples are set forth to aid in the understanding of the invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.

Unless otherwise indicated, <sup>1</sup>H NMRs were run on a Bruker AC-300.

20

#### EXAMPLE 1

##### 1-(3,4-Methylenedioxyphenyl)-2-[5-(3-trifluoromethylphenyl)furoyl]-2,3,4,9-tetrahydro-1H-β-carboline (#58)

To a suspension of 5-(3-trifluoromethylphenyl)furoic acid (256 mg, 1 mmol) in DCM (20 mL, anhydrous) was added oxalyl chloride (165 mg, 1.3 mmol), followed by two drops of DMF. The mixture was stirred at room temperature for 1h. A solution of 1-(3,4-methylenedioxyphenyl)-2,3,4,9-tetrahydro-1H-β-carboline (292 mg, 1 mmol) (prepared according to the process as disclosed in WO97/43287, Intermediate 7, page 24) and triethylamine (0.4 mL) in DCM (10 mL, anhydrous) was added and the mixture was stirred at room temperature for 16h, washed sequentially with aqueous NaHCO<sub>3</sub>, brine (2X), 1N HCl and brine (2X) and dried with MgSO<sub>4</sub>. After evaporation of the solvent, a white solid was obtained.

30

mp: 126-129 °C

MS (m/z): 531 (MH<sup>+</sup>);

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 2.96 (d, *J* = 8 Hz, 1 H), 3.24 (m, 1 H), 3.56 (m, 1 H), 4.60 (d, *J* = 8 Hz, 1 H), 5.90 (s, 2 H), 6.70 (d, *J* = 8 Hz, 1 H), 6.83-6.99 (m, 4 H), 7.13-7.34 (m, 4 H), 7.55 (m, 3 H), 7.87 (d, *J* = 7 Hz, 1 H), 7.95 (s, 1 H), 8.23 (s, 1 H);

5

## EXAMPLE 2

### 9-[2-(Pyrrolidin-1-yl)-ethyl]-1-(3,4-methylenedioxyphenyl)-2-[5-(3-trifluoromethyl-phenyl)furoyl]-2,3,4-trihydro-1H-β-carboline (#75)

10 To a solution of 1-(3,4-methylenedioxyphenyl)-2-[5-(3-trifluoromethyl phenyl)furoyl]-2,3,4,9-tetrahydro-1H-β-carboline (prepared as in Example 1) (600 mg, 1.14 mmol) in DMF (15 mL, anhydrous) was added sodium hydride (60%, 105 mg, 2.6 mmol) at room temperature. The mixture was stirred at room temperature for 30 min. N-chloroethylpyrrolidine hydrochloride (214 mg, 1.26 mmol) and 15-crown  
15 ether-5 (1 drop) were added. The mixture was stirred at room temperature for 16h, quenched with NH<sub>4</sub>Cl, extracted with ethyl acetate and dried with MgSO<sub>4</sub>. After evaporation of the solvent, the residue was purified by column chromatography (silica gel, ethyl acetate:hexanes = 3:1) to yield a white solid.

MS (m/z): 628 (MH<sup>+</sup>)

20 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.26 (m, 4 H), 2.64 (m, 4 H), 2.89 (m, 2 H), 3.05 (d, *J* = 8 Hz, 1 H), 3.28 (t, *J* = 8 Hz, 1 H), 3.59 (t, *J* = 8 Hz, 1 H), 3.96 (m, 1 H), 4.16 (m, 1 H), 4.58 (d, *J* = 8 Hz, 1 H), 5.96 (s, 2 H), 6.75 (d, *J* = 8 Hz, 1 H), 6.84 (m, 2 H), 7.02 (d, *J* = 8 Hz, 1 H), 7.15-7.29 (m, 4 H), 7.43 (s, 1 H), 7.59 (m, 3 H), 7.89 (d, *J* = 7 Hz, 1 H), 7.96 (s, 1 H)

25

The corresponding methanesulfonic acid salt was prepared by addition of 1.0 equivalent of methanesulfonic acid to a solution of the title compound in DCM to produce product for biological testing.

mp: 122-124°C.

30

### EXAMPLE 3

#### 1-(3,4-Methylenedioxyphenyl)-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#7)

A solution of 1-(3,4-methylenedioxyphenyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline  
5 (3.73 g, 12.8 mmol) (prepared according to the process as disclosed in  
WO97/43287, Intermediate 7, page 24) and 2-chloro-5-(3,4-  
dimethoxyphenyl)pyrimidine (1.6 g, 6.4 mmol) in DMF (50 mL, anhydrous) was  
heated at 120°C under stirring for 16h. The reaction mixture was quenched with  
NH<sub>4</sub>Cl and extracted with ethyl acetate. The organic phase was washed with brine  
10 (2X) and dried with MgSO<sub>4</sub>. Column chromatography (silica gel, ethyl  
acetate:hexanes = 2:3) yielded a white solid.

mp: 173-175 °C

MS (m/z): 507 (MH<sup>+</sup>)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)  $\delta$  2.91 (d, *J* = 9 Hz, 1 H), 3.02 (td, *J* = 9, 1 Hz, 1 H), 3.39 (td,  
15 *J* = 9, 1 Hz, 1 H), 3.92 (s, 3 H), 3.94 (s, 3 H), 5.02 (d, *J* = 9, 1 Hz, 1 H), 5.92 (s, 2 H),  
6.72 (d, *J* = 8 Hz, 1 H), 6.87-7.03 (m, 4 H), 7.11-7.17 (m, 3 H), 7.31 (d, *J* = 8 Hz, 1  
H), 7.56 (d, *J* = 8 Hz, 1 H), 7.80 (s, 1 H), 8.56 (s, 2 H)

20

### EXAMPLE 4

#### 1-(3,4-Methylenedioxyphenyl)-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]-9- dimethylaminoethyl-2,3,4-trihydro-1H- $\beta$ -carboline (#5)

Following the procedure outlined in Example 2, 1-(3,4-  
methylenedioxyphenyl)-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]-2,3,4,9-tetrahydro-  
25 1H- $\beta$ -carboline (prepared as in Example 3) (1.0 g, 1.97 mmol), 2-chloro-N,N-  
dimethylethylamine hydrochloride (0.342 g, 2.37 mmol), sodium hydride (60%, 0.190  
g, 4.74 mmol) and 15-crown ether-5 were reacted to yield the product as a slightly  
yellow solid (after column chromatography with silica gel, ethyl acetate).

MS (m/z): 578 (MH<sup>+</sup>)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)  $\delta$  2.21 (s, 6 H), 2.22 (m, 1 H), 2.61 (m, 1 H), 2.89 (dd, *J* =  
30 13, 4 Hz, 1 H), 3.03 (td, *J* = 13, 4 Hz, 1 H), 3.35 (td, *J* = 13, 4 Hz, 1 H), 3.91 (m, 1 H),

3.92 (s, 3 H), 3.95 (s, 3 H), 4.06 (m, 1 H), 4.96 (dd,  $J = 13, 4$  Hz, 1 H), 5.93 (s, 2 H), 6.72 (d,  $J = 8$  Hz, 1 H), 6.83 (d, 1 H), 6.85-6.98 (m, 4 H), 7.12 (d,  $J = 8$  Hz, 1 H), 7.21 (d,  $J = 8$  Hz, 1 H), 7.31 (d,  $J = 8$  Hz, 1 H), 7.34 (s, 1 H), 7.58 (d,  $J = 8$  Hz, 1 H), 8.56 (s, 2 H)

5

### Example 5

1-(3,4-Methylenedioxyphenyl)-2-[5-(4-methoxyphenyl)-pyrimidin-2-yl]-4-oxo-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#157) &

10 1-(3,4-Methylenedioxy-phenyl)-2-[5-(4-methoxyphenyl)-pyrimidin-2-yl]-4-hydroxy-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#158)

To a mixture of DDQ (113.5 mg, 0.5 mmol) and 1-(3,4-methylenedioxyphenyl)-2-[5-(3,4-dimethoxyphenyl)pyrimidin-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (prepared as in Example 3) (51 mg, 0.1 mmol) was added a mixed solvent of THF:water (9:1) at -78°C. The mixture was stirred at 0°C and allowed to warm to room temperature over a period of 15 h. Column chromatography (silica gel, hexanes:ethyl acetate = 1:1) yielded the-oxo- and -hydroxy derivatives respectively, as white solids.

#### #157:

20 MS (m/z) 521 ( $MH^+$ ), 519 ( $M-1$ )

$^1H$  NMR ( $CDCl_3$ )  $\delta$  3.90 (d,  $J = 18$  Hz, 1 H), 3.89 (s, 3 H), 3.91 (s, 3 H), 5.43 (d,  $J = 18$  Hz, 1 H), 5.84 (s, 2 H), 6.62 (d,  $J = 8$  Hz, 1 H), 6.71 (d,  $J = 8$  Hz, 1 H), 6.88-7.00 (m, 4 H), 7.29-7.43 (m, 3 H), 7.53 (s, 1 H), 8.25 (m, 1 H), 8.51 (s, 2 H); 9.55 (s, 1 H)

25 #158:

MS (m/z) 523 ( $MH^+$ ), 521 ( $M-1$ )

$^1H$  NMR ( $CDCl_3$ )  $\delta$  3.30 (t,  $J = 6$  Hz, 1 H), 3.69 (d,  $J = 6$  Hz, 1 H), 3.92 (s, 3 H), 3.94 (s, 3 H), 5.97 (s, 2 H), 6.11 (s, 1 H), 6.71 (d,  $J = 8$  Hz, 1 H), 6.93-7.05 (m, 4 H), 7.18 (d,  $J = 8$  Hz, 1 H), 7.23 (d,  $J = 8$  Hz, 1 H), 7.40 (t,  $J = 6$  Hz, 1 H), 7.49 (d,  $J = 8$  Hz, 1 H), 7.82 (d,  $J = 8$  Hz, 1 H), 8.43 (s, 2 H); 9.15 (s, 1 H)

30

## EXAMPLE 6

1-(3,4-Methylenedioxyphenyl)-2-[4-(4-methoxyphenyl)thiazol-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#169)

- 5 A. 1-(3,4-Methylenedioxyphenyl)-2-[3-(fluorenylmethyloxycarbonyl)thiocarbamoyl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline

A mixture of 1-(3,4-methylenedioxyphenyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (2.66 g, 9.08 mmol) (prepared according to the process as disclosed in WO97/43287, Intermediate 7, page 24) and Fmoc-isothiocyanate (2.82 g, 10.14  
10 mmol) was dissolved in dry dichloromethane (50 mL). The mixture was stirred for 16 hours at ambient temperature, and then concentrated *in vacuo*. Purification by flash chromatography (0-10% methanol in dichloromethane) yielded the protected thiourea as a pale yellow solid.

MS (m/z): 574 (MH<sup>+</sup>)

- 15 <sup>1</sup>H-NMR (CDCl<sub>3</sub>)  $\delta$  2.86 (dd, *J* = 12.9, 5.1 Hz, 1 H), 3.09 (dt, *J* = 17.1, 6.9 Hz, 1 H), 3.56 (dt, *J* = 12.9, 5.1 Hz, 1 H), 4.19 (t, *J* = 6.9 Hz, 1 H), 4.43-4.53 (m, 2 H), 5.91 (s, 2 H), 6.70 (d, *J* = 8 Hz, 1 H), 6.90 (br d, *J* = 7.6 Hz, 1 H), 6.97 (br s, 1 H), 7.11-7.78 (series of m, 17 H)

- 20 B. 1-(3,4-Methylenedioxyphenyl)-2-(thiocarbamoyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline

A solution of the protected thiourea from Part A (4.78 g, 8.33 mmol) in 20% (v/v) piperidine in methanol was heated to reflux for 5 h. The mixture was concentrated *in vacuo* to yield a crude residue which was purified by flash  
25 chromatography (SiO<sub>2</sub>, 0-10% methanol in dichloromethane) to yield a yellow solid.

MS (m/z): 352 (MH<sup>+</sup>)

- <sup>1</sup>H-NMR (CDCl<sub>3</sub>)  $\delta$  2.69-2.87 (series of m, 2 H), 3.10-3.19 (m, 1 H), 4.24 (br s, 1 H), 6.00 (d, *J* = 3.3 Hz, 2 H), 6.72 (d, *J* = 8.0 Hz, 1 H), 6.87 (d, *J* = 8.0 Hz,  
30 1 H), 7.00-7.11 (series of m, 3 H), 7.30 (d, *J* = 8.0 Hz, 1H), 7.46 (d, *J* = 7.7 Hz, 1H), 7.74 (br s, 3 H), 11.06 (s, 1 H)

C. 1-(3,4-Methylenedioxyphenyl)-2-[4-(4-methoxyphenyl)thiazol-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#169)

To a solution of the thiourea from Part B (223 mg, 0.63 mmol) in a 1:1  
5 mixture of dioxane:ethanol (5 mL) was added 4-methoxyphenyl-2'-  
bromoacetophenone (175 mg, 0.76 mmol) and triethylamine (0.40 mL). The  
mixture was heated to 70°C for 3 h, cooled to room temperature and  
concentrated in a rotary evaporator. The residue was purified by flash  
chromatography (SiO<sub>2</sub>, 0-10% methanol in dichloromethane) to yield a colorless  
10 solid.

MS (m/z): 482 (MH<sup>+</sup>)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)  $\delta$  2.86-2-3.07 (series of m, 2 H), 3.61-3.71 (m, 1 H), 3.78  
(s, 3 H), 3.91-4.02 (m, 1 H), 5.99 (d, *J* = 3.3 Hz, 2 H), 6.58 (s, 1 H), 6.80-7.11  
(series of m, 8 H), 7.31 (d, *J* = 7.8 Hz, 1 H), 7.48 (d, *J* = 7.6 Hz, 1 H), 7.82 (d, *J* =  
15 8.7 Hz, 2 H), 10.93 (s, 1 H)

#### EXAMPLE 7

1-(3,4-Methylenedioxyphenyl)-2-[4-phenylthiazol-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -  
20 carboline (#170)

A. 1-(3,4-Methylenedioxyphenyl)-2-[3-(fluorenylmethyloxycarbonyl)  
thiocarbamoyl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline

A mixture of 1-(3,4-methylenedioxyphenyl)-2,3,4,9-tetrahydro-1H- $\beta$ -  
carboline (2.66 g, 9.08 mmol) (prepared according to the process as disclosed in  
25 WO97/43287, Intermediate 7, page 24) and Fmoc-isothiocyanate (2.82 g, 10.14  
mmol) was dissolved in dry dichloromethane (50 mL). The mixture was stirred for  
16 hours at ambient temperature, and then concentrated *in vacuo*. Purification  
by flash chromatography (0-10% methanol in dichloromethane) yielded the  
protected thiourea as a pale yellow solid.

30 MS (m/z): 574 (MH<sup>+</sup>)

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  2.86 (dd,  $J = 12.9, 5.1$  Hz, 1 H), 3.09 (dt,  $J = 17.1, 6.9$  Hz, 1 H), 3.56 (dt,  $J = 12.9, 5.1$  Hz, 1 H), 4.19 (t,  $J = 6.9$  Hz, 1 H), 4.43-4.53 (m, 2 H), 5.91 (s, 2 H), 6.70 (d,  $J = 8$  Hz, 1 H), 6.90 (br d,  $J = 7.6$  Hz, 1 H), 6.97 (br s, 1 H), 7.11-7.78 (series of m, 17 H)

5

B. 1-(3,4-Methylenedioxyphenyl)-2-(thiocarbamoyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline

A solution of the protected thiourea from Part A (4.78 g, 8.33 mmol) in 20% (v/v) piperidine in methanol was heated to reflux for 5 h. The mixture was concentrated *in vacuo* to yield a crude residue which was purified by flash chromatography ( $\text{SiO}_2$ , 0-10% methanol in dichloromethane) to yield a yellow solid.

MS ( $m/z$ ): 352 ( $\text{MH}^+$ )

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  2.69-2.87 (series of m, 2 H), 3.10-3.19 (m, 1 H), 4.24 (br s, 1 H), 6.00 (d,  $J = 3.3$  Hz, 2 H), 6.72 (d,  $J = 8.0$  Hz, 1 H), 6.87 (d,  $J = 8.0$  Hz, 1 H), 7.00-7.11 (series of m, 3 H), 7.30 (d,  $J = 8.0$  Hz, 1H), 7.46 (d,  $J = 7.7$  Hz, 1H), 7.74 (br s, 3 H), 11.06 (s, 1 H)

C. 1-(3,4-Methylenedioxyphenyl)-2-[4-phenylthiazol-2yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#170)

To a solution of the thiourea of Part B (227 mg, 0.65 mmol) was added  $\beta$ -bromoacetophenone (159 mg, 0.80 mmol) and triethylamine (0.40 mL). This mixture was heated to  $70^\circ\text{C}$  for 3 h, cooled to room temperature and concentrated in a rotary evaporator. The residue was purified by flash chromatography ( $\text{SiO}_2$ , 0-10% methanol in dichloromethane) to yield a pale yellow solid.

MS ( $m/z$ ): 452 ( $\text{MH}^+$ )

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$  2.87-2-3.06 (series of m, 2 H), 3.63-3.73 (m, 1 H), 3.93-3.99 (m, 1 H), 5.99 (d,  $J = 3.3$  Hz, 2 H), 6.59 (s, 1 H), 6.81-7.11 (series of m, 5 H), 7.25-7.69 (series of m, 6 H), 7.89 (d,  $J = 7.4$  Hz, 2 H), 10.95 (s, 1 H)



## EXAMPLE 8

1-(2,3-Dihydro-benzofuran-5-yl)-2-[5-(2,3-dimethyl-3H-imidazol-4-yl)-pyrimidin-2-yl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#190)

5

2-(5-bromo-2-pyrimidinyl)-1-(2,3-dihydro-5-benzofuranyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (0.45 g, 1.00 mmol), 1,2-dimethyl-1H-imidazole (0.18 g, 1.87 mmol), Pd(OAc)<sub>2</sub> (12 mg, 0.05 mmol), PPh<sub>3</sub> (26 mg, 0.1 mmol) and K<sub>2</sub>CO<sub>3</sub> (0.28 g, 2 mmol) were stirred in 3.5 mL DMF at 140°C for 14 hours. The mixture was poured into aqueous 10% NaOH solution (50 mL). The resulting solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3×50 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Purification by preparative TLC yielded the title product as yellow powder.

10

<sup>1</sup>H NMR 300 MHz (CDCl<sub>3</sub>)  $\delta$  2.21 (s, 3H), 2.35 (s, 3H), 2.90 (m, 2H), 3.10 (t, 2H, *J* = 8.8.Hz), 3.35 (m, 1H), 4.52 (t, 2H, *J* = 8.8.Hz), 4.91 (m, 1H), 6.68 ~ 7.61 (m, 10 H)

15

MS (m/z) 463 (MH<sup>+</sup>), 461 (MH<sup>-</sup>).

## EXAMPLE 9

20 2-[2,3']Bipyridinyl-6'-yl-1-(2,3-dihydro-benzofuran-5-yl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (#191)

A: 2-(5-Bromo-pyridin-2-yl)-1-(2,3-dihydro-benzofuran-5-yl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline

25

1-(2,3-dihydro-5-benzofuranyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline (11.6 g, 40 mmol), 2, 5-dibromopyridine (10.42 g, 44 mmol), Pd<sub>2</sub>dba<sub>3</sub> (1.465 g, 1.6 mmol), dppp (1.32 g, 3.2 mmol) and NaOtBu (5.38 g, 56 mmol) were stirred in 60 mL DMF at 80°C for 3 days. The reaction mixture was filtered through a plug of Celite with CH<sub>2</sub>Cl<sub>2</sub>. The reaction mixture was then concentrated, the crude mixture was then loaded on Foxy column (110 g silica gel) and eluted with ethyl acetate / hexane (3:7). The product crystallized out in test tubes. The product

30

was concentrated and then recrystallized from THF to yield the product as yellow crystals.

$^1\text{H}$  NMR 400 MHz (THF- $d_8$ )  $\delta$  0.91 (m, 1H), 1.15 (m, 1H), 1.25(t, 2H,  $J$  = 9.5 Hz), 1.60 (m, 1H), 2.31 (m, 1H), 2.60 (t, 2H,  $J$  = 9.5 Hz), 4.75 (d, 1H,  $J$  = 7.6 H), 5.02 (d, 1H,  $J$  = 7.6 Hz), 5.10 ~5.28 (m, 4H), 5.380 (m, 2H), 5.58 (m, 1H), 5.72 (m, 1H), 6.28 (s, 1H), 8.12 (s, 1H)

MS (m/z) 446, 448 ( $\text{MH}^+$ ), 444, 446 ( $\text{MH}^-$ ).

B: 2-[2,3']Bipyridinyl-6'-yl-1-(2,3-dihydro-benzofuran-5-yl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline


The product from step A above, (0.4 g, 0.896 mmol), 2-tributylstannanyl-pyridine (0.8 g, 2.17 mmol) and  $\text{Pd}(\text{PPh}_3)_4$  (0.12 g, 0.104 mmol) were stirred in 1,4-dioxane (5 mL) at 88°C for 24 h. The reaction mixture was filtered through a plug of Celite with  $\text{CH}_2\text{Cl}_2$  and then concentrated to a small volume. Preparative TLC (3:7 ethyl acetate / hexane; then 5%  $\text{CH}_3\text{OH}$  /  $\text{CH}_2\text{Cl}_2$ ) yielded the product as a yellow solid.


$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.82 (m, 1H), 3.10 (m, 3H), 3.58 (m, 1H), 4.31 (m, 1H), 4.53 (t, 2H,  $J$  = 9.5 Hz), 6.71 (d, 1H,  $J$  = 7.6 Hz), 6.85 (d, 1H,  $J$  = 7.6 Hz)

MS (m/z) 445, ( $\text{MH}^+$ ), 443 ( $\text{MH}^-$ )

Following procedures as described herein, the compounds as listed in Tables 1-6 were prepared.

TABLE 1




Cmpd #	R <sup>D</sup>	R <sup>2</sup>		R <sup>3</sup>
1	H	3,4-methylene dioxyphe <sup>n</sup> yl	-	-
2	dimethyl aminoethyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-benzothiazolyl	-
3	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-benzothiazolyl	-
4	dimethyl aminoethyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(4-chlorophenyl)
5	dimethyl aminoethyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(3,4-dimethoxyphenyl)
6	dimethyl aminoethyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(4-methoxyphenyl)
7	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(3,4-dimethoxyphenyl)
8	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(2-nitro-4-methylsulfonylphenyl)
9	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(4-methoxyphenyl)
10	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(4-chlorophenyl)
148	2-(N-pyrrolidinyl) ethyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-pyrimidinyl	5-(4-methoxyphenyl)
150	2-benzoxazolyl	3,4-methylene dioxyphe <sup>n</sup> yl	2-benzoxazolyl	-
151	H	3,4-methylene dioxyphe <sup>n</sup> yl	2-benzoxazolyl	-

153	H	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(2-pyridyl)
154	2-(N- pyrrolidinyl) ethyl	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(2-nitro-4- methylsulfonyl phenyl)
155	dimethyl aminoethyl	3,4-methylene dioxyphenyl	2-benzoxazolyl	-
156	dimethyl aminoethyl	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(2-pyridyl)
161	H	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(4-methyl phenyl)
166	5-(4-methyl phenyl)-2- pyrimidinyl	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(4-methyl phenyl)
167	H	3,4-dimethoxy phenyl	2-pyrimidinyl	5-(3,4-dimethoxy phenyl)
168	H	3,4-methylene dioxyphenyl	2-pyrimidinyl	-
169	H	3,4-methylene dioxyphenyl	2-thiazolyl	4-(4-methoxy phenyl)
170	H	3,4-methylene dioxyphenyl	2-thiazolyl	4-phenyl
172	H	3,4-methylene dioxyphenyl	2-benz imidazolyl	-
173	H	3,4-methylene dioxyphenyl	2-pyrimidinyl	-
174	H	5-(2,3-dihydro)- benzofuryl	2-pyrimidinyl	5-(4-methoxy phenyl)
175	H	3,4-methylene dioxyphenyl	2-pyrimidinyl	5-(3,4-dimethoxy phenyl)

176	H	R-3,4-methylene dioxypheyl	2-pyrimidinyl	5-(3,4-dimethoxy phenyl)
177	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(4-hydroxy phenyl)
178	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(4-[2-(N-pyrrolidinyl)ethoxy]phenyl)
179	H	3,4-methylene dioxypheyl	2-pyrimidinyl	5-bromo
180	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(4-(2-(4-morpholinyl)ethoxy)phenyl)
181	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(4-(2-(dimethyl amino)ethoxy) phenyl)
182	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(2-pyridyl)
183	H	3,4-methylene dioxypheyl	2-pyrimidinyl	5-(2-pyridyl)
184	H	6-(2,3-dihydrobenzo-[1,4]-dioxin-6-yl)	2-pyrimidinyl	5-(4-methoxy phenyl)
185	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(3-pyridyl)
186	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-bromo
187	H	5-(2,3-dihydro)-benzofuryl	2-pyrimidinyl	5-(2-pyridyl)

188	H	6-(2,3-dihydrobenzo-[1,4]-dioxin-6-yl)	2-pyrimidinyl	5-(4-methoxyphenyl)
189	H	3,4-methylene dioxyphehyl	2-pyrimidinyl	5-(2-pyridyl)
190	H	5-(2,3-dihydro benzofuryl)	2-pyrimidinyl	5-(2,3-dimethyl-3H-imidazol-4-yl)
191	H	5-(2,3-dihydro benzofuryl)	2-pyridyl	5-(2-pyridyl)

TABLE 2



Cmpd #	R <sup>D</sup>	R <sup>2</sup>	Y	C	R <sup>3</sup>
11	H	3,4-methylene dioxyphehyl	CH <sub>2</sub>	2-furyl	5-(2-chloro-5-trifluoromethylphenyl)
12	H	3,4-methylene dioxyphehyl	CH <sub>2</sub>	2-furyl	5-(3-trifluoromethylphenyl)
13	dimethyl aminoethyl	3,4-methylene dioxyphehyl	C(O)	2-benzo(b)furyl	5-nitro
14	H	3,4-methylene dioxyphehyl	C(O)	2-benzo(b)furyl	5-nitro
15	H	3,4-methylene	C(O)	2-	6-methoxy

		dioxyphenyl		benzo(b)furyl	
16	H	3,4-methylene dioxyphenyl	C(O)	2- benzo(b)furyl	-
17	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-benzo(b) thienyl	-
18	H	3,4-methylene dioxyphenyl	C(O)	2-benzo(b) thienyl	-
19	2-(3-nitro phenyl)-5- furylcarbonyl	3,4-(difluoro methylene dioxy)phenyl	C(O)	2-furyl	5-(3-nitro phenyl)
20	2-(3-trifluoro methylphenyl )5-furyl carbonyl	3,4-(difluoro methylene dioxy)phenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
21	H	3,4-(difluoro methylene dioxy)phenyl	C(O)	2-furyl	5-(3-nitro phenyl)
22	H	3,4-(difluoro methylene dioxy)phenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
23	2-methoxy carbonylethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
24	2-methoxy carbonyl methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-nitro phenyl)
25	2-methoxy carbonyl methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
26	2-pyridyl methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
27	4-chloro-n-	3,4-methylene	C(O)	2-furyl	5-(3-trifluoro

	butyl	dioxyphenyl			methylphenyl)
28	4-morpholinyl ethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
29	carboxyethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
30	carboxy methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-nitro phenyl)
31	carboxy methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
33	diethylamino ethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
34	dimethyl aminobutyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
35	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
36	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5- <i>t</i> -butyl
37	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-nitro phenyl)
38	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-nitro phenyl)
39	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-chloro phenyl)
40	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- chlorophenyl)
41	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro phenyl)
42	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- methylphenyl)
43	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-chloro-5- trifluoromethyl



					phenyl)
44	dimethyl aminopropyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
74	4-pyridyl methyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
75	pyrrolidinyl ethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
45	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-amino phenyl)
46	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-chloro phenyl)
47	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- chlorophenyl)
48	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-[4-(3- carboxy)-n- propylcarbonyl aminophenyl]
49	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-acetyl aminophenyl)
50	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(4-nitro phenyl)
51	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- chloro phenyl)
52	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-[4-(3- carboxy)-n- propylcarbonyl aminophenyl]
53	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- methylphenyl)
54	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-nitro phenyl)

55	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-acetyl aminophenyl)
56	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro phenyl)
57	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-chloro-5- trifluoromethyl phenyl)
58	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
59	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(3-amino phenyl)
60	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-[4-(4-hydroxy -n-butyl)amino phenyl]
61	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-[2-(dimethyl amino)methyl carbonylamino phenyl]
62	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-trifluoro methyl
63	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-bromo
64	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-nitro
65	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5- <i>t</i> -butyl
66	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-(2-nitro-4- chlorophenyl)
78	H	3-pyridyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
79	H	4-	C(O)	2-furyl	5-(3-trifluoro

		chlorophenyl			methylphenyl)
80	H	4-cyanophenyl	C(O)	2-furyl	5-(3-nitro phenyl)
81	H	4-cyanophenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
82	H	4-dimethyl aminophenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
83	H	4-dimethyl aminophenyl	C(O)	2-furyl	5-(3-nitro phenyl)
84	H	4-nitrophenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
85	H	4-nitrophenyl	C(O)	2-furyl	5-(3-nitro phenyl)
86	H	4-pyridyl	C(O)	2-furyl	5-(3-nitro phenyl)
87	H	4-pyridyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
88	H	phenyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
89	H	3,4-methylene dioxyphenyl	C(O)	2-thiazolyl	4-methyl-5-(4-trifluoro methylphenyl)
90	H	3,4-methylene dioxyphenyl	C(O)	2-thienyl	4-phenyl-5-trifluoromethyl
91	H	3,4-methylene dioxyphenyl	C(O)	2-thienyl	5-(4-chloro phenyl)
92	H	3,4-dimethyl phenyl	C(O)	3-(1,2,5-triazolyl)	1-phenyl-4-methyl
93	H	3,4-dichloro phenyl	C(O)	3-(1,2,5-triazolyl)	1-phenyl-4-methyl
94	H	3,4-dimethoxy	C(O)	3-(1,2,5-	1-phenyl-4-

		phenyl		triazolyl)	methyl
95	H	3,4-dimethyl phenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
96	H	3,4-methylene dioxyphenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
97	H	3,4-methylene dioxyphenyl	C(O)	3-(1,2,5- triazolyl)	5-(3-pyridyl)
98	H	3-trifluoro methyl-4- chlorophenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
99	H	4-cyano phenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
100	H	4-methoxy carbonyl phenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
101	H	4-methoxy phenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
102	H	4-nitrophenyl	C(O)	3-(1,2,5- triazolyl)	1-phenyl-4- methyl
103	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	3-furyl	2-methyl-5-(4- chloro phenyl)
104	H	3,4-methylene dioxyphenyl	C(O)	3-furyl	2-methyl-5- phenyl
105	H	3,4-methylene dioxyphenyl	C(O)	3-furyl	2-trifluoro methyl-5-(4- chlorophenyl)
106	H	3,4-methylene dioxyphenyl	C(O)	3-pyrazolyl	1-phenyl-5- methyl
107	H	4-[N-(3-di methylamino)- n-propoxy]	C(O)	3-pyrazolyl	1-phenyl-5- methyl


		phenyl			
108	H	3,4-methylene dioxyphenyl	C(O)	3-pyridyl	6-chloro
109	H	3,4-dichloro phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
110	H	3,4-dimethoxy phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
111	H	3,4-dimethyl phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
112	H	3,4-methylene dioxyphenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
113	H	3,5-dimethyl phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
114	H	3-trifluoro methyl-4- chlorophenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
115	H	4- cyanophenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
116	H	4-methoxy carbonyl phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
117	H	4-methoxy phenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
118	H	4-nitrophenyl	C(O)	4-isoxazolyl	3-phenyl-5- methyl
119	H	3,4-dimethyl phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
120	H	3,4-dichloro phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
121	H	3,4-dimethoxy phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl

122	H	3,4-methylene dioxyphenyl	C(O)	4-pyrazolyl	1-phenyl-5-n- propyl
123	H	3,4-methylene dioxyphenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
124	H	3,4-methylene dioxyphenyl	C(O)	4-pyrazolyl	1-(4-chloro phenyl)-5- trifluoromethyl
125	H	3,4-methylene dioxyphenyl	C(O)	4-pyrazolyl	1-(4- nitrophenyl)-5- trifluoromethyl
126	H	3,5-dimethyl phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
127	H	3-trifluoro methyl-4- chlorophenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
128	H	4- cyanophenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
129	H	4-methoxy carbonyl phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
130	H	4-methoxy phenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
131	H	4-nitrophenyl	C(O)	4-pyrazolyl	1-phenyl-5- trifluoromethyl
132	H	3,4-methylene dioxyphenyl	C(O)	4-thiazolyl	2-(4-pyrazinyl)
133	H	3,4-dichloro phenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
134	H	3,4-dimethoxy phenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
135	H	3,4-dimethyl	C(O)	5-pyrazolyl	1-benzyl-3-t-

		phenyl			butyl
136	H	3,4-methylene dioxyphenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
137	H	3,5-dimethyl phenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
138	H	3-trifluoro methyl-4- chlorophenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
139	H	4- cyanophenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
140	H	4-methoxy carbonyl phenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
141	H	4-methoxy phenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
142	H	4-nitrophenyl	C(O)	5-pyrazolyl	1-benzyl-3-t- butyl
143	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	naphthyl	-
144	H	3,4-methylene dioxyphenyl	C(O)	naphthyl	-
145	H	3,4-methylene dioxyphenyl	SO <sub>2</sub>	2-thienyl	5-phenyl sulfonyl
146	H	3,4-methylene dioxyphenyl	SO <sub>2</sub>	2-thienyl	3-phenyl sulfonyl
147	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-benzofuryl	-
149	H	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-phenyl
152	dimethyl aminoethyl	3,4-methylene dioxyphenyl	C(O)	2-furyl	5-phenyl

185	H	3,4-methylene dioxyphenyl	CH <sub>2</sub>	phenyl	-
-----	---	------------------------------	-----------------	--------	---

TABLE 3





Cmpd #	R <sup>2</sup>	R <sup>4</sup>	Y		R <sup>3</sup>
67	3,4-methylene dioxyphenyl	carboxy	C(O)	2-furyl	5-(3-nitro phenyl)
68	3,4-methylene dioxyphenyl	2-(dimethyl amino)ethoxy carbonyl	C(O)	2-furyl	5-(3-nitro phenyl)
69	3,4-methylene dioxyphenyl	2-(dimethyl amino)ethyl aminocarbonyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
70	3,4-methylene dioxyphenyl	carboxy	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
71	3,4-methylene dioxyphenyl	methoxy carbonyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
72	3,4-methylene dioxyphenyl	methoxy carbonyl	C(O)	2-furyl	5-(3-trifluoro methylphenyl)
73	3,4-methylene dioxyphenyl	methoxy carbonyl	C(O)	2-furyl	5-(3-nitro phenyl)



TABLE 4






Cmpd #	R <sup>2</sup>	Y		R <sup>3</sup>
76	3,4-methylene dioxypheyl	C(O)	2-furyl	5-(3-nitrophenyl)
77	3,4-methylene dioxypheyl	C(O)	2-furyl	5-(4-chlorophenyl)

TABLE 5



5

Cmpd #	R <sup>2</sup>	Z		R <sup>3</sup>
157	3,4-methylene dioxypheyl	C(O)	2-pyrimidinyl	5-(3,4-dimethoxy phenyl)
158	3,4-methylene dioxypheyl	CHOH	2-pyrimidinyl	5-(3,4-dimethoxy phenyl)
159	3,4-methylene dioxypheyl	C(O)	2-pyrimidinyl	5-(4-methylphenyl)
160	3,4-methylene dioxypheyl	CHOH	2-pyrimidinyl	5-(4-methylphenyl)

162	3,4-methylene dioxyphenyl	C(O)	2-pyrimidinyl	5-(4-methoxyphenyl)
163	3,4-methylene dioxyphenyl	C(O)	2-pyrimidinyl	5-(4-methoxyphenyl)
164	3,4-methylene dioxyphenyl	CHOH	2-pyrimidinyl	5-(4-methoxyphenyl)
165	3,4-methylene dioxyphenyl	C(O)	2-pyrimidinyl	5-(4-methoxyphenyl)

### EXAMPLE 10

#### IN VITRO TESTING

##### 5 Cyclic Nucleotide Phosphodiesterase (PDE) Assay

##### PDEV Isolation

PDEV was isolated from rabbit and human tissues according to the protocol described by Boolell et al. (*Boolell, M., Allen, M. J., Ballard, S. A., Gelfo-Attee, S., Muirhead, G. J., Naylor, A. M., Osterloh, I. H., and Gingell, C*) in *International*

10 *Journal of Impotence Research* **1996** 8, 47-52 with minor modifications.

Briefly, rabbit or human tissues were homogenized in an ice-cold buffer solution containing 20mM HEPES (pH 7.2), 0.25M sucrose, 1mM EDTA, and 1mM phenylmethylsulphonyl fluoride (PMSF). The homogenates were centrifuged at  
 15 100,000g for 60 minutes at 4°C. The supernatant was filtered through 0.2µM filter and loaded on a Pharmacia Mono Q anion exchange column (1ml bed volume) that was equilibrated with 20mM HEPES, 1mM EDTA and 0.5mM PMSF. After washing out unbound proteins, the enzymes were eluted with a linear gradient of 100-600 mM NaCl in the same buffer (35 to 50 ml total, depending on the tissue. Enzymes  
 20 from the skeletal muscle, corpus cavernosum, retina, heart and platelet were eluted with 35, 40, 45, 50, and 50 ml respectively. ) The column was run at a flow rate of 1ml/min and 1ml fractions were collected. The fractions comprising various PDE activities were pooled separately and used in later studies.

### Measurement of Inhibition of PDEV

The PDE assay was carried out as described by Thompson and Appleman in *Biochemistry* **1971** 10, 311-316 with minor modifications, as noted below.

5        The assays were adapted to a 96-well format. The enzyme was assayed in 5mM MgCl<sub>2</sub>, 15mM Tris HCl (pH 7.4), 0.5 mg/ml bovine serum albumin, 1 μM cGMP or cAMP, 0.1 μCi [<sup>3</sup>H]-cGMP or [<sup>3</sup>H]-cAMP, and 2-10 μl of column elution. The total volume of the assay was 100 μl. The reaction mixture was incubated at 30°C for 30 minutes. The reaction was stopped by boiling for 1 minute and then cooled down on  
10 ice. The resulting [<sup>3</sup>H] 5'-mononucleotides were further converted to uncharged [<sup>3</sup>H]-nucleosides by adding 25 μl 1 mg/ml snake venom (*Ophiophagus hannah*) and incubating at 30°C for 10 minute. The reaction was stopped by the addition of 1 ml Bio-Rad AG1-X2 resin slurry (1:3). All the charged nucleotides were bound by the resin and only the uncharged [<sup>3</sup>H]-nucleosides remained in the supernatant after  
15 centrifuging. An aliquot of 200 μl was taken and counted by liquid scintillation. PDE activity was expressed as pmol cyclic nucleotide hydrolyzed/min/ml of enzyme preparation.

Inhibitor studies were carried out in assay buffer with a final concentration of  
20 10% DMSO. Under these conditions, the hydrolysis of product increased with time and enzyme concentration in a linear fashion.

### Example 11

#### 25        In Vitro Determination of K<sub>i</sub> for Phosphodiesterase Inhibitors:

The assay was adapted to a 96-well format. Phosphodiesterase was assayed in 5mM MgCl<sub>2</sub>, 15mM Tris HCl (pH 7.4), 0.5 mg/ml bovine serum albumin, 30 nM <sup>3</sup>H-cGMP and test compound at various concentrations. The amount of  
30 enzyme used for each reaction was such that less than 15% of the initial substrate was converted during the assay period. For all measurements, the test compound

was dissolved and diluted in 100% DMSO (2%DMSO in assay). The total volume of the assay was 100  $\mu$ l. The reaction mixture was incubated at 30°C for 90 minutes. The reaction was stopped by boiling for 1 minute and then immediately cooled by transfer to an ice bath. To each well was then added 25  $\mu$ l 1 mg/ml snake venom  
 5 (*Ophiophagus hannah*) and the reaction mixture incubating at 30°C for 10 minute. The reaction was stopped by the addition of 1 ml Bio-Rad AG1-X2 resin slurry (1:3). An aliquot of 200  $\mu$ l was taken and counted by liquid scintillation.

The % inhibition of the maximum substrate conversion (by the enzyme in  
 10 the absence of inhibitor) was calculated for each test compound concentration. Using GraphPad Prism's nonlinear regression analysis (sigmoidal dose response), the % inhibition vs log of the test compound concentration was plotted to determine the IC<sub>50</sub>. Under conditions where substrate concentration  $\ll$  K<sub>m</sub> of the enzyme (K<sub>m</sub> = substrate concentration at which half of the maximal velocity of  
 15 the enzyme is achieved), K<sub>i</sub> is equivalent to the IC<sub>50</sub> value.

Mass spec and PDEV inhibitory activity for representative compounds of the present invention are described in Tables 6-7. Inhibitory data is presented either as  
 20 the IC<sub>50</sub> ( $\mu$ M), as a percent inhibition at a given concentration of test compound or as a K<sub>i</sub> value.

TABLE 6

Cmpd #	MW	MS (M+1)	%Inh @ 10 $\mu$ M (rabbit)
1	288.30	289	67
2	496.63	497	43
3	425.51	426	84
4	552.08	553	94
5	577.68	578	

6	547.66	548	95
7	506.56	507	91
8	569.60	580	89
9	476.53	477	80
10	480.95	481	79
11	550.96	551	53
12	516.52	517	76
13	552.58	553	94
14	481.46	482	88
15	466.49	467	90
16	436.47	437	88
17	523.65	524	88
18	452.53	453	87
19	758.64	759	15
20	804.64	805	4
21	543.48	544	32
22	566.48	567	16
23	616.59	617	30
24	579.56	580	40
25	602.56	603	20
26	621.61	622	88
27	621.05	622	56
28	643.66	644	94
29	602.56	601* (M-1) No M+1 ion	79
30	565.54	564	93
31	588.54	587* (M-1) no M+1 ion	75
33	629.68	630	88
33	629.68	630	63
34	629.68	630	67

35	601.62	602	98
36	513.63	514	97
37	578.62	579	95
38	578.62	579	91
39	568.07	569	93
40	613.07	614	78
41	578.62	579	93
42	592.65	593	89
43	636.07	637	22
44	615.65	616	65
45	477.52	478	
46	496.95	497	
47	541.94	542	83
48	577.59	578	60
49	519.55	520	60
50	507.50	508	76
51	541.94	542	
52	577.59	578	76
53	521.53	522	85
54	507.50	508	81
55	519.55	520	70
56	507.50	508	
57	564.95	565	76
58	530.50	531	
59	477.52	478	92
60	549.62	550	89
61	562.62	563	90
62	454.40	455	86
63	465.30	465	78
64	431.40	432	83
65	442.51	443	66

66	541.94	542	23
67	550.50	549* (M-1) No M+1 ion	50
68	622.63	623	35
69	644.65	645	21
70	574.51	573	46
71	588.54	587* (M-1) No M+1 ion	21
72	588.54	587* (M-1) No M+1 ion	15
73	565.54	564* (M-1) No M+1 ion	20
74	621.61	622	84
75	627.66	628	
76	524.55	525	27
77	514.00	515	58
78	487.48	488	35
79	520.94	521	37
80	488.50	489	27
81	511.50	510* (M-1) No M+1 ion	18
82	529.56	530	13
83	506.56	507	27
84	531.49	532	20
85	508.49	509	26
86	464.48	465	69
87	487.48	488	34
88	486.49	487	49
89	561.58	562	
90	546.57	547	69
91	513.01	514	82

92	461.57	462	39
93	502.40	503	44
94	493.56	494	19
95	461.57	462	13
96	477.52	478	57
97	480.55	481	71
98	535.95	536	38
99	458.52	459	40
100	491.55	492	24
101	463.54	464	48
102	478.51	479	40
103	510.97	511	55
104	476.53	477	72
105	564.95	565	40
106	476.53	477	70
107	533.67	534	15
108	431.88	432	48
109	502.40	503	31
110	493.56	494	32
111	461.56	462	35
112	477.52	478	33
113	461.56	462	29
114	535.95	536	27
115	458.52	459	30
116	491.54	492	32
117	463.53	464	32
118	478.51	479	28
119	514.55	515	28
120	555.39	556	18
121	546.55	547	10
122	504.59	505	65



123	530.50	531	56
124	564.95	565	53
125	575.50	576	54
126	514.55	515	12
127	588.94	589	13
128	511.51	512	11
129	544.53	545	46
130	516.52	517	45
131	531.49	532	12
132	480.55	481	76
133	557.52	556	1
134	548.68	547	5
135	516.69	517	8
136	532.64	533	18
137	516.69	517	-3
138	591.07	592	13
139	513.64	514	-9
140	546.67	547	8
141	518.66	519	11
142	533.63	534	-5
143	517.63	518	60
144	446.50	447	76
145	578.69	579* (M-1) No M+1 ion	43
146	578.69	579	35

TABLE 7

Cmpd #	MW	MS (M+1)	IC <sub>50</sub> (μM)	% INH @10 μM (rabbit)
147	507.59	508	5.2	52
148	573.69	574	4.4	66

149	462.50	463	0.75	80
150	526.55	527		4
151	409.44	410	0.13	95
152	533.63	534	1.2	80
153	447.50	448	0.12	95
154	666.76	667	0.11	97
155	480.56	481	9.2	57
156	518.62	519	0.0075	
157	520.54	521	0.0087 <sup>a</sup>	
158	522.56	523		49
159	474.52	475	0.024	
160	476.53	477	8.95	
161	460.54	461	0.789	
162	490.52	491	0.024 <sup>a</sup>	
164	492.53	493		82
166	628.73	629		58
167	522.60	523	1.49	
168	370.41	371	2.15 <sup>a</sup>	
169	481.57	482	0.042	
170	451.55	452	0.049	
172	408.46	409		44 <sup>a</sup>
174	474.56	475	0.150 <sup>a</sup>	
175	506.56	507	0.214	
176	506.56	507	0.056	
178	557.70	558		37
179	449.31	450	1.58	
180	573.69	574		40
182	445.52	446	0.058	
183	447.50	448	0.122 <sup>a</sup>	
184			0.640 <sup>a</sup>	
185		446		200.40 <sup>a</sup>

186		448		240.10 <sup>a</sup>
187		446		49.79 <sup>a</sup>
188		491		642.69 <sup>a</sup>
189		448		121.60 <sup>a</sup>
190		463		Ki = 14.21 nM
191		443		Ki = 0.69 nM

<sup>a</sup> Compounds tested using human tissue.

#### Example 12

5

#### IN VIVO TESTING

Following the procedure disclosed by Carter et al., (Carter, A. J., Ballard, S. A., and Naylor, A. M.) in The Journal of Urology 1998, 160, 242-246, compounds of the present invention are tested for in vivo efficacy.

10

#### Example 13

As a specific embodiment of an oral composition, 100 mg of the compound of Example 7 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

15


While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the

20

following claims and their equivalents.

We claim:

1. A compound of the formula (I):



5 wherein

$R^1$  is independently selected from the group consisting of halogen, nitro, hydroxy,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy,  $-NH_2$ ,  $-NHR^A$ ,  $-N(R^A)_2$ ,  $-OR^A$ ,  $-C(O)NH_2$ ,  $-C(O)NHR^A$ ,  $-C(O)N(R^A)_2$ ,  $-NC(O)-R^A$ ,  $-SO_2NHR^A$ ,  $-SO_2N(R^A)_2$ , phenyl (optionally substituted with 1 to 3  $R^B$ ) and heteroaryl (optionally substituted with 1 to 3  $R^B$ );

10 where each  $R^A$  is independently is independently selected from the group consisting of  $C_1$ - $C_8$ alkyl, aryl (optionally substituted with 1 to 3  $R^B$ ),  $C_1$ - $C_8$ aralkyl (optionally substituted with 1 to 3  $R^B$ ) and heteroaryl (optionally substituted with 1 to 3  $R^B$ );

15 where each  $R^B$  is independently selected from the group consisting of halogen, hydroxy, nitro, cyano,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy,  $C_1$ - $C_8$ alkoxycarbonyl, carboxy $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkylsulfonyl, trifluoromethyl, trifluoromethoxy, amino, acetylamino, di( $C_1$ - $C_8$ alkyl)amino, di( $C_1$ - $C_8$ alkyl)amino $C_1$ - $C_8$ alkoxy, di( $C_1$ - $C_8$ alkyl)aminoacetyl $C_1$ - $C_8$ alkyl, di( $C_1$ - $C_8$ alkyl)aminoacetylamino, carboxy $C_1$ - $C_8$ alkylcarbonylamino, hydroxy $C_1$ - $C_8$ alkylamino,  $NHR^A$ ,  $N(R^A)_2$  and  
20 heterocycloalkyl $C_1$ - $C_8$ alkoxy;

$n$  is an integer from 0 to 4;

$X$  is selected from the group consisting of O, S and  $NR^D$ ;

25 where  $R^D$  is selected from the group consisting of hydrogen, hydroxy,  $-OR^A$ ,  $C_1$ - $C_8$ alkyl (wherein the alkyl is optionally substituted with one to three substituent independently selected from halogen, carboxy, amino,  $C_1$ - $C_8$ alkylamino, di( $C_1$ - $C_8$ alkyl)amino,  $C_1$ - $C_8$ alkoxycarbonyl, heteroaryl or heterocycloalkyl), heteroaryl and

heteroarylcarbonyl (wherein the heteroaryl may be optionally substituted with phenyl or substituted phenyl, where the phenyl substituents are one to three  $R^B$ );

$R^2$  is selected from the group consisting of  $C_5$ - $C_{10}$ alkyl (optionally substituted with 1 to 3  $R^C$ ), aryl (optionally substituted with 1 to 3  $R^B$ ), heteroaryl (optionally substituted with 1 to 3  $R^B$ ) and heterocycloalkyl (optionally substituted with 1 to 3  $R^B$ );

where each  $R^C$  is independently selected from the group consisting of halogen, hydroxy, nitro,  $NH_2$ ,  $NHR^A$  and  $N(R^A)_2$ ;


$Z$  is selected from the group consisting of  $CH_2$ ,  $CHOH$  and  $C(O)$ ; provided that when  $Z$  is  $CHOH$  or  $C(O)$ , then  $X$  is  $NH$ ;

$R^4$  is selected from the group consisting of hydrogen, hydroxy, carboxy,  $C_1$ - $C_6$ alkylcarbonyl,  $C_1$ - $C_6$ alkoxycarbonyl,  $di(C_1$ - $C_8$ alkyl)aminoalkoxycarbonyl,  $di(C_1$ - $C_8$ alkyl)amino $C_1$ - $C_8$ alkylaminocarbonyl, and  $-COR^F$ ;

where  $R^F$  is selected from the group consisting of  $C_1$ - $C_8$ alkyl,  $NH_2$ ,  $NHR^A$ ,  $NR^A_2$ ,  $-C_1$ - $C_8$ alkyl- $NH_2$ ,  $-C_1$ - $C_8$ alkyl- $NHR^A$ ,  $-C_1$ - $C_8$ alkyl- $NR^A_2$  and  $-NH$ - $C_1$ - $C_8$ alkyl- $NR^A_2$ ;

$a$  is an integer from 0 to 1;


$Y$  is selected from the group consisting of  $CH_2$ ,  $C(O)$ ,  $C(O)O$ ,  $C(O)-NH$  and  $SO_2$ ;



is selected from the group consisting of naphthyl, heteroaryl and heterocycloalkyl;

$m$  is an integer from 0 to 2;

$R^3$  is independently selected from the group consisting of halogen, nitro,  $C_1$ - $C_8$ alkyl,  $C_1$ - $C_8$ alkoxy, trifluoromethyl, trifluoromethoxy, phenyl (optionally substituted with 1 to 3  $R^B$ ), phenylsulfonyl, naphthyl,  $C_1$ - $C_8$ alkyl, heteroaryl (optionally substituted with 1 to 3  $R^B$ ),  $NH_2$ ,  $NHR^A$  and  $N(R^A)_2$ ;



provided that when is 2-furyl or 2-thienyl, then  $m$  is an integer from 1 to 2;

and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1 wherein

n is 0;


5 X is selected from S or  $\text{NR}^{\text{D}}$ , wherein  $\text{R}^{\text{D}}$  is selected from the group consisting of hydrogen, halo $\text{C}_1\text{-C}_6$ alkyl, di( $\text{C}_1\text{-C}_4$ alkyl)amino $\text{C}_1\text{-C}_6$ alkyl, heteroaryl, heteroaryl $\text{C}_1\text{-C}_4$ alkyl, heterocycloalkyl $\text{C}_1\text{-C}_4$ alkyl, carboxy $\text{C}_1\text{-C}_4$ alkyl,  $\text{C}_1\text{-C}_4$ alkoxycarbonyl $\text{C}_1\text{-C}_4$ alkyl and heteroarylcarbonyl; wherein the heteroaryl is further optionally substituted with phenyl or substituted phenyl, wherein the substituents on the phenyl  
10 are one to two independently selected from  $\text{R}^{\text{B}}$ ;

where each  $\text{R}^{\text{B}}$  is independently selected from the group consisting of halogen, nitro,  $\text{C}_1\text{-C}_4$ alkyl,  $\text{C}_1\text{-C}_4$ alkoxy, trifluoromethyl, trifluoromethoxy, amino and di( $\text{C}_1\text{-C}_4$ alkyl)amino;

$\text{R}^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl, 3,4-  
15 (difluoro)methylenedioxyphenyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzo-[1,4]-dioxin-6-yl, pyridyl, phenyl and substituted phenyl; wherein the phenyl substituents are one to two substituents independently selected from halogen,  $\text{C}_1\text{-C}_4$ alkyl,  $\text{C}_1\text{-C}_4$ alkoxy, trifluoromethyl, cyano, nitro,  $\text{C}_1\text{-C}_4$ alkoxycarbonyl, di( $\text{C}_1\text{-C}_4$ alkyl)amino or di( $\text{C}_1\text{-C}_4$ alkyl)amino $\text{C}_1\text{-C}_4$ alkoxy;

20  $\text{R}^4$  is selected from the group consisting of hydrogen, carboxy,  $\text{C}_1\text{-C}_4$ alkoxycarbonyl, di( $\text{C}_1\text{-C}_4$ alkyl)amino $\text{C}_1\text{-C}_4$ alkoxycarbonyl and di( $\text{C}_1\text{-C}_4$ alkyl)amino $\text{C}_1\text{-C}_4$ alkylaminocarbonyl;


Y is selected from the group consisting of  $\text{C}(\text{O})$ ,  $\text{SO}_2$  and  $\text{CH}_2$ ;




is selected from the group consisting of naphthyl and heteroaryl;

25  $\text{R}^3$  is independently selected from the group consisting of halogen, nitro,  $\text{C}_1\text{-C}_4$ alkyl,  $\text{C}_1\text{-C}_4$ alkoxy, trifluoromethyl,  $\text{C}_1\text{-C}_4$ alkyl, pyrazinyl, pyridyl, halogen substituted pyridyl, dimethyl substituted imidazolyl, phenyl, phenylsulfonyl and substituted phenyl; wherein the substituents on the phenyl are one or more substituents independently selected from halogen, hydroxy,  $\text{C}_1\text{-C}_4$ alkyl,  $\text{C}_1\text{-C}_4$ alkoxy,  
30 trifluoromethyl, trifluoromethoxy, nitro, amino, acetamino,  $\text{C}_1\text{-C}_4$ alkylsulfonyl,

carboxyC<sub>1</sub>-C<sub>4</sub>alkylcarbonylamino, hydroxyC<sub>1</sub>-C<sub>4</sub>alkylamino, di(C<sub>1</sub>-C<sub>4</sub>alkyl)aminoC<sub>1</sub>-C<sub>4</sub>alkoxy, di(C<sub>1</sub>-C<sub>4</sub>alkyl)aminoacetyl amino or heterocycloalkylC<sub>1</sub>-C<sub>4</sub>alkoxy;



provided that when  is 2-furyl or 2-thienyl, then m is an integer from 1 to 2;


5 and pharmaceutically acceptable salts thereof.


3. The compound of Claim 2, wherein

X is selected from S or NR<sup>D</sup>, where R<sup>D</sup> is selected from the group consisting of hydrogen, di(methyl)aminoethyl, di(methyl)amino-n-propyl, di(ethyl)aminoethyl,  
10 di(ethyl)amino-n-butyl, N-pyrrolidinylethyl, N-morpholinylethyl, 2-pyridylmethyl, 4-pyridylmethyl, 5-(4-methylphenyl)-2-pyrimidinyl, carboxymethyl, carboxyethyl, 4-chloro-n-butyl, 2-(5-(3-trifluoromethylphenyl)furyl)carbonyl, 2-(5-(3-nitrophenyl)furyl)carbonyl, methoxycarbonylmethyl, methoxycarbonylethyl and 2-benzoxazolyl;


15 R<sup>2</sup> is selected from the group consisting of phenyl, 3,4-methylenedioxyphenyl, 3,4-(difluoro)methylenedioxyphenyl, 2,3-dihydrobenzofuryl, 2,3-dihydrobenzo-[1,4]-dioxin-6-yl, 4-pyridyl, 3-pyridyl, 4-cyanophenyl, 3-nitrophenyl, 4-nitrophenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4-dimethylphenyl, 3,5-dimethylphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethyl-4-chlorophenyl, 3,4-dichlorophenyl, 4-chlorophenyl, 4-methoxycarbonylphenyl, 3,4-dimethoxyphenyl, 4-(dimethylamino)phenyl and 4-(N-(3-dimethylamino)-n-propoxy)phenyl;


R<sup>4</sup> is selected from the group consisting of hydrogen, carboxy, dimethylaminoethoxycarbonyl, dimethylaminoethylaminocarbonyl and methoxycarbonyl;



25  is selected from the group consisting of naphthyl, 2-pyrimidinyl, 2-furyl, 3-furyl, 2-benzofuryl, 2-thienyl, 2-benzothienyl, 2-benzothiazolyl, 2-benzoxazolyl, 2-benzimidazolyl, 4-thiazolyl, 2-thiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 3-(1,2,5-triazolyl), 4-isoxazolyl, 2-pyridyl and 3-pyridyl;

$R^3$  is independently selected from the group consisting of chloro, bromo, methyl, n-propyl, t-butyl, methoxy, trifluoromethyl, nitro, phenyl, benzyl, phenylsulfonyl, 4-hydroxyphenyl, 4-chlorophenyl, 4-methylphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 5-trifluoromethylphenyl, 4-methoxyphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 4-aminophenyl, 2-nitro-4-chlorophenyl, 2-nitro-4-methylphenyl, 2-nitro-4-methylsulfonylphenyl, 3-acetylaminophenyl, 4-acetylaminophenyl, 4-(3-carboxy-n-propyl)carbonylaminophenyl, 2-chloro-5-trifluoromethylphenyl, 4-(4-hydroxy-n-butyl)aminophenyl, 2-(dimethylamino)acetylaminophenyl, 4-[2-(N-pyrrolidinyl)ethoxy]phenyl, 4-[2-(4-morpholinyl)ethoxy]phenyl, 4-(2-(dimethylamino)ethoxy)phenyl, 4-pyrazinyl, 2,3-dimethyl-3H-imidazolyl, 2-pyridyl and 3-pyridyl;



provided that when  is 2-furyl or 2-thienyl, then m is an integer from 1 to 2;

and pharmaceutically acceptable salts thereof.


4. The compound of Claim 3 wherein


X is  $NR^D$ , where  $R^D$  is selected from the group consisting of hydrogen, di(methyl)aminoethyl, 4-pyridylmethyl, 2-pyridylmethyl, N-morpholinylethyl, carboxyethyl, carboxymethyl, di(ethyl)aminoethyl, N-pyrrolidinylethyl and 5-(4-methylphenyl)-2-pyrimidinyl;

$R^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl, 2,3-dihydrobenzofuryl and 2,3-dihydrobenzo-[1,4]-dioxin-6-yl;

$R^4$  is hydrogen;

Y is selected from the group consisting of C(O) and  $CH_2$ ;



 is selected from the group consisting of naphthyl, 2-pyrimidinyl, 2-furyl, 2-benzofuryl, 2-thienyl, 2-benzothieryl, 2-benzothiazolyl, 2-benzoxazolyl, 2-thiazolyl, 4-thiazolyl and 2-pyridyl;



m is an integer from 0 to 1;

$R^3$  is selected from the group consisting of bromo, t-butyl, methoxy, trifluoromethyl, nitro, phenyl, 4-chlorophenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 4-methylphenyl, 4-methoxyphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 2-nitro-4-chlorophenyl, 2-nitro-4-methylphenyl, 2-nitro-4-methylsulfonylphenyl, 4-(3-carboxy-n-propyl)carbonylaminophenyl, 2-chloro-5-trifluoromethylphenyl, 4-(4-hydroxy-n-butyl)aminophenyl, 2-2-(dimethylamino)acetylaminophenyl, 4-pyrazinyl, 2-pyridyl and 2,3-dimethyl-3H-imidazol-4-yl;

10 provided that when  is 2-furyl or 2-thienyl, then m is 1; and pharmaceutically acceptable salts thereof.


5. The compound of Claim 4 wherein

X is  $NR^D$ , where  $R^D$  is selected from the group consisting of hydrogen, di(methyl)aminoethyl, N-morpholinylethyl, carboxymethyl and N-pyrrolidinylethyl;


$R^2$  is selected from the group consisting of 3,4-methylenedioxyphenyl and 2,3-dihydrobenzofuryl;

Z is selected from the group consisting of  $CH_2$  and  $C(O)$ ; provided that when Z is  $C(O)$ , then X is NH;

20 Y is  $C(O)$ ;

 is selected from the group consisting of 2-pyrimidinyl, 2-furyl, 2-benzofuryl, 2-benzoxazolyl, 2-thiazolyl and 2-pyridyl;

$R^3$  is selected from the group consisting of t-butyl, methoxy, nitro, phenyl, 4-chlorophenyl, 4-methylphenyl, 4-methoxyphenyl, 3,4-dimethoxyphenyl, 3-trifluoromethylphenyl, 2-nitrophenyl, 3-nitrophenyl, 4-nitrophenyl, 3-aminophenyl, 2-nitro-4-methylsulfonylphenyl, 2-(dimethylamino)acetylaminophenyl, 2-pyridyl and 2,3-dimethyl-3H-imidazol-4-yl;



provided that when is 2-furyl, then m is 1;  
and pharmaceutically acceptable salts thereof.

6. A compound of Claim 3 selected from the group consisting of
  - 1-(3,4-methylenedioxyphenyl)-2-[(5-phenyl-2-furyl)carbonyl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline;
  - 1-(3,4-methylenedioxyphenyl)-2-[5-(2-pyridyl)-2-pyrimidinyl]-9-di(methyl)aminoethyl-2,3,4,9-tetrahydro-1H- $\beta$ -carboline;
  - 1-(3,4-methylenedioxyphenyl)-2-[5-(3,4-dimethoxyphenyl)-2-pyrimidinyl]-1,2,3,9-tetrahydro-4-oxo-4H- $\beta$ -carboline
  - 1-(3,4-methylenedioxyphenyl)-2-[5-(4-methylphenyl)-2-pyrimidinyl]-1,2,3,9-tetrahydro-4-oxo-4H- $\beta$ -carboline;
  - 1-(3,4-methylenedioxyphenyl)-2-[5-(4-methoxyphenyl)-2-pyrimidinyl]-1,2,3,4-tetrahydro-4-oxo-4H- $\beta$ -carboline;
  - 1-(3,4-methylenedioxyphenyl)-2-[4-(4-methoxyphenyl)-2-thiazolyl]-2,3,4,9-tetrahydro-1H- $\beta$ -carboline;
  - 1-(3,4-methylenedioxyphenyl)-2-(4-phenyl-2-thiazolyl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline;
  - 2-[2,3']Bipyridinyl-6'-yl-1-(2,3-dihydro-benzofuran-5-yl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline;
  - 1-(2,3-dihydrobenzofuran-5-yl)-2-[5-(2,3-dimethyl-3H-imidazol-4-yl)-2,3,4,9-tetrahydro-1H- $\beta$ -carboline]; and pharmaceutically acceptable salts thereof.
7. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claim 1.
8. A pharmaceutical composition made by mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.

9. A process for making a pharmaceutical composition comprising mixing a compound of Claim 1 and a pharmaceutically acceptable carrier.
- 5 10. A method of treating sexual dysfunction in a subject in need thereof comprising administering to the subject a therapeutically effective amount of the compound of Claim 1.
11. The method of treating sexual dysfunction of Claim 10, wherein the sexual  
10 dysfunction is male erectile dysfunction.
12. The method of Claim 10, wherein the sexual dysfunction is selected from the group consisting of male sexual dysfunction, male erectile dysfunction, impotence, female sexual dysfunction, female sexual arousal dysfunction and  
15 female sexual dysfunction related to blood flow and nitric oxide production in the tissues of the vagina and clitoris.
13. A method for increasing the concentration of cGMP in penile tissue in a male subject in need thereof comprising administering to the subject an effective amount  
20 of the compound of Claim 1.
14. A method of treating a condition selected from the group consisting of male erectile dysfunction (ED), impotence, female sexual arousal dysfunction, female sexual dysfunction related to blood flow and nitric oxide production in the tissues of  
25 the vagina and clitoris, premature labor, dysmenorrhea, cardiovascular disorders, atherosclerosis, arterial occlusive disorders, thrombosis, coronary rest stenosis, angina pectoris, myocardial infarction, heart failure, ischemic heart disorders, hypertension, pulmonary hypertension, asthma, intermittent claudication and diabetic complications in a subject in need thereof comprising administering to the  
30 subject a therapeutically effective amount of the compound of Claim 1.